

# Mineral Contents of Several Indonesian Rice Varieties

*by Susiyanti Susiyanti*

---

**Submission date:** 30-Jan-2023 06:03AM (UTC+0700)

**Submission ID:** 2001813234

**File name:** 12.\_Susiyanti\_mineral\_content.pdf (1,000.29K)

**Word count:** 2855

**Character count:** 14350

# Mineral Contents of Several Indonesian Rice Varieties

\*Susiyanti Susiyanti, Nurmayulis Nurmayulis, and  
Sulastris Isminingsih

Department of Agroecotechnology, Agriculture Faculty,  
University of Sultan Agung Tirtayasa, Serang, Banten Province,  
Indonesia

\*susiyanti@untirta.ac.id

Rahmayety Rahmayety and Yeyen Maryani

Department of Chemical Engineering, Engineering  
Faculty, University of Sultan Agung Tirtayasa, Serang,  
Banten Province, Indonesia

Rahmayety99@gmail.com

Suseno Amin

Department of Agroecotechnology, Agriculture Faculty  
University of Padjajaran, West Jawa Province, Indonesia

Susenoamin73@gmail.com

**Abstract** – Rice is an important food commodity in the world, even the majority of Indonesia's population consumes rice as a staple food. The chemical composition of rice consumed is very complex. Rice contains organic ingredients and minerals. At present, the mineral as a nutrient has not been widely recognized by the public as well as the adequacy of calories, protein or vitamins. Analysis of mineral content in several Indonesian rice varieties (such as Kewal Bulu Hideung, Jalahawara, Pandan Wangi, Rojolele, Bendang Pulau, Batang Piaman, Cisantana, Sokan, Ciherang, Sidrap, and Kewal Gudril) carried out in Shimadzu laboratory, Osaka-Japan, using EDX-7000/8000 Energy-Dispersive X-ray Fluorescence Spectrometers. With these tools a variety of macro and micro minerals can be analyzed from each rice sample, including phosphorus (P), potassium (K), sulfur (S), calcium (Ca), manganese (Mn), copper (Cu), zinc (Zn), iron (Fe), Rubidium (Rb), silicon (Si). Based on the results, the mineral content for various types of kinds of rice tested was varied. The highest mineral content was found in Kewal Bulu Hideung rice (3.884%), and the lowest was Sokan (1.677%). The mineral content found in succession from the largest to the smallest found in all rice tested were P, K, S, Ca, and Zn. The mineral content found only in special varieties were Mn, Fe, Rb, Si, Cu.

**Keywords** : rice, variety, Indonesia, mineral content

## I. INTRODUCTION

Rice is an important food commodity in the world, even most of the population of Indonesia consume rice as a staple food. The chemical composition of rice consumed is very complex. In addition to carbohydrate, rice has a mineral content that serves to other growth and development of the vital organs of a human being. According to [1], more than half of the world population is our suffering from bioavailable nutrient deficiencies particularly mainly in developing countries. Many things affect the mineral content contained in rice, such as the environment, genetic factors and process of rice miller and the other on food processing [2].

The mineral has an important function to human health, such as forming tissues of the body, regulate, and control the metabolic processes in the body. The mineral can be grouped

into macrominerals (Ca, P, Mg, Na, K, Cl, S) and micro minerals (Fe, Zn, Cu, I, Mn, Cr, Co, Se, Mo, F). The demand for minerals that can be supplied through food consumption.

In Indonesia, rice, besides having a role in contributing to the adequacy of protein, energy, also plays a role in fulfilling minerals such as iron. Seeing these conditions, encourage plant breeders to assemble superior rice varieties that have good mineral content for health. For example, Bengawan Solo (high Ca), Limboto (P content) Cimelati (Fe content), and many others.

This mineral content is very needed by the body, but if found in excess content in food will cause a serious problem for health. Recently, heavy metal poisoning from foodstuffs has increased [3]. According to [4], heavy metal contamination in humans can also through food and water consumed. Environmental pollution by heavy metals can occur if industries that use these metals do not pay attention to environmental safety, especially when disposing of their waste. Hazardous metals in high concentrations will be very dangerous if found in water, soil and air environments [5]. Furthermore, all plants that grow on the contaminated soil will accumulate these metals in all parts (roots, stems, leaves, and fruit). Food produced from polluted environments will cause accumulation of contaminants into their organs.

Considering the position of rice which is considered as the world's important food and the main food of the Indonesian people, it is very necessary to inform the mineral content contained in the food they consume. Based on these problems, this study wants to examine the content of minerals in rice in some local Indonesian rice.

## II. MATERIALS AND METHODS

Analysis of mineral content in several superior and local Indonesian rice varieties (such as Kewal Bulu Hideung, Jalahawara, Pandan Wangi, Rojolele, Bendang Pulau, Batang Piaman, Cisantana, Sokan, Ciherang, Sidrap, and Kewal

Gudril) was carried out at Shimadzu Laboratory, Osaka - Japan, using EDX-7000/8000 Energy-Dispersive X-ray Fluorescence Spectrometers. The EDX-7000/8000 incorporates a new high-performance semiconductor detector, provides excellent sensitivity, resolution, and throughput for an array of applications, from general screening analysis to advanced materials research in such fields as chemicals. With these tools various macro and micro minerals can be analyzed from each rice sample, including iron (Fe), manganese (Mn), copper (Cu), zinc (Zn), calcium (Ca), magnesium (Mg), sodium (Na), potassium (K), phosphorus (P), sulfur (S), boron (B), molybdenum (Mo), Cu, nickel (Ni), and others.

III. RESULTS AND DISCUSSION

The rice used is a type of rice originating from several regions such as Banten area (Kewal Bulu Hideung, Jalahawara, Kewal Gudril), West Java (such as: Ciherang, Pandan Wangi, Rojolele), West Sumatra (Bendang Pulau, Batang Piaman, Sokan), Sulawesi (Cisantana, Sidrap). The physical appearance of the rice tested can be seen in Figure 1. From its appearance, the rice from Sidrap has a slender form, while the other rice is categorized based on the ratio of length per width as medium rice.

Based on the results of the analysis using EDX, the mineral content of P, K, S, Ca, Mn, Cu, Zn, Fe, Rb, Si is obtained (can be seen in Table 1 & 2). The complete profile of the mineral content found in the Kewal Bulu Hideung rice can be seen in Figure 2; Jalahawara (Figure 3), Pandan Wangi (Figure 4), Rojolele (Figure 5), Bendang Pulau (Figure 6), Batang Piaman (Figure 7), Cisantana (Figure 8), Sokan (Figure 9), Ciherang (Figure 10), Sidrap (Figure 11), and Kewal Gudril (Figure 12).




Figure 1. Physical appearance of rice samples: Kewal Bulu Hideung (A1), Jalahawara (A2), Pandan Wangi (A3), Rojolele (A4), Bendang Pulau (A5), Batang Piaman (A6), Cisantana (A7), Sokan (A8), Ciherang (A9), Sidrap (A10), and Kewal Gudril (A11).

TABLE 1. MINERAL CONTENT OF P, K, S, Ca, Mn, Cu ON RICE SOURCED FROM SEVERAL INDONESIAN RICE VARIETIES

No.	Jenis Padi (%)	P	K	S	Ca	Mn	Cu
1	Kewal Bulu Hideung	2,157	1,246	0,196	0,001	0,004	0,001
2	Jalahawara	1,126	0,393	0,381	0,031		
3	Pandan Wangi	1,525	0,431	0,274	0,014	0,003	
4	Rojolele	1,361	0,39	0,254	0,01		
5	Bendang Pulau	1,131	0,3	0,229	0,008		
6	Batang Piaman	1,034	0,221	0,309	0,019		
7	Cisantana	1,823	0,54	0,271	0,025	0,003	
8	Sokan	0,973	0,228	0,329	0,016		
9	Ciherang	1,565	0,405	0,221	0,019		
10	Sidrap	1,862	0,597	0,307	0,016		
11	Kewal Gudril	1,288	0,827	0,209	0,063	0,007	

The mineral content for various rice is varied (Table 1 & 2, Figures 2 to 12). The highest mineral content was found in Kewal Bulu Hideung rice (3.884%), and the lowest was Sokan (1.677%). The mineral content found in all rice varieties is P, K, S, Ca, and Zn. The mineral content found only in special varieties is Mn, Fe, Rb, Si, Cu. The main minerals in rice are Ca Cl, Na, Si, Fe, P, and K, followed by Si and Mg [6]. The rice in this study contained Ca, Zn, Mn, Fe, Rb, Si, and Cu which were low, while the P content was proportionally higher. The distribution of minerals in rice seeds is mostly found in the outer layer of the seeds and further downward. Many things affect the mineral content in plant organs, such as genetic factors, soil and water environment where the rice grows, and post-harvest processing.

TABLE 2. MINERAL CONTENT of Zn, Fe, Rb, Si, ORGANIC MATERIAL, TOTAL MINERALS ON RICE SOURCED from SEVERAL INDONESIAN RICE VARIETIES

No.	Jenis Padi (%)	Zn	Fe	Rb	Si	BO	Mineral
1	Kewal Bulu Hideung	0,006	0,006	0,005	0,264	96,116	3,884
2	Jalahawara	0,003	0,002			98,063	1,937
3	Pandan Wangi	0,003	0,003	0,002	0,016	97,578	2,422
4	Rojolele	0,003		0,002		97,98	2,02
5	Bendang Pulau	0,002		0,003	0,113	98,213	1,787
6	Batang Piaman	0,003		0,003	0,189	98,223	1,777
7	Cisantana	0,003	0,002	0,002	0,126	97,204	2,796
8	Sokan	0,003			0,0128	98,323	1,677

9	Ciherang	0.003				97.787	2.213
10	Sidrap	0.003	0.003			97.213	2.787
11	Kewal Gudril	0.004	0.004	0.003	1.043	96.561	3.439




Figure 2. Profile of mineral content of Kewal Bulu Hideung rice




Figure 3. Profile of mineral content of Jalawahara rice




Figure 4. Profile of mineral content of Pandan Wangi rice




Figure 5. Profile of mineral content of Gambar Rojolele rice




Figure 6. Profile of mineral content of Bendang Pulau rice




Figure 7. Profile of mineral content of Batang Piaman rice




Figure 8. Profile of mineral content of Cisantana rice




Figure 9. Profile of mineral content of Sokan rice




Figure 10. Profile of mineral content of Ciherang rice




Figure 11. Profile of mineral content of Sidrap rice




Figure 12. Profile of mineral content of Kewal Gudril rice

The results showed a range of P, K and Ca contents respectively 0.973-2.157% for P; 0.221-1.246% for K; and 0.001-0.382% for Ca. Based on research, [7] reported rice mineral content in Nigerian rice varieties, such as P content (0.52-0.54%); K (0.15-0.20%); Ca (0.09-0.11%). This means that Indonesian rice that has been studied has a higher P and K content, while its Ca content is lower. According to [8], P, K, and Ca accumulated around the aleurone layer, and most widely found on the lateral side of the rice seed.

The results of the study in Table 1 show that the Cu content is 0.001% (only found in Kewal Bulu Hideung), while the Zn content ranges from 0.002-0.006% (Table 2). The content of Cu, and Zn in aromatic and non-aromatic rice in India, found that the Cu content ranged from 0.0041-0.01595%; and Zn 0.0093-0.017% [9]. When compared, the content of Cu and Zn was found in all samples of rice varieties tested in this study, apparently lower than the rice studied in India. Zn and Cu were found on localization different from P, because the Zn and Cu were the most abundant in endosperm in the closed position layer (i.e. the layer of sub-aleurone), and spread deeper into the endosperm [8].

The Fe content of the analysis ranges from 0.002-0.006% and is only found in Kewal Bulu Hideung, Jalahawara, Pandan Wangi, Cisantana, Sidrap, Kewal Gudril. The Fe content in these varieties is generally higher when compared to the Fe content in rice studied by [9], which is 0.0008-0.0031%. The localization sites of the aleurone layer and the inner starch endosperm. The pattern of distribution of Fe abounds in the dorsal side, where some vascular bundles, present and acting as a gateway for translocation [8].

Mn content in this study ranged from 0.005-0.007% found in several varieties, namely Kewal Bulu Hideung, Pandan Wangi, Cisantana and Kewal Gudril. In the body, Mn acts as a catalyst for several important metabolic reactions in proteins, carbohydrates, and fats. In protein metabolism, Mn activates the conversion of amino acids with specific enzymes

such as arginase, prolinase, dipeptidase. In carbohydrate metabolism, Mn plays an active role in several conversion reactions in glucose oxidation and synthesis of oligosaccharides. In fat metabolism, Mn acts as a cofactor in the synthesis of long chain fatty acids and cholesterol.

The content of a metal content in plant organs will also be affected by the availability of other metals. According to [10], the hull of high accumulation of the elements, such as N and P from the hull into the brown rice. The levels of Si ranged from 0.016-1.043% found in the Kewal Bulu Hideung rice, Pandan Wangi, Bendang Pulau, Batang Piaman, Cisantana, and Kewal Gudril.

In this study, Rb content of 0.002-0.005% was found in Kewal Bulu Hideung rice, Pandan Wangi, Rojolele, Bendang Pulau, Cisantana, and Kewal Gudril. The Rb concentration in white grain ranged from 1.2-5.5 mg kg<sup>-1</sup> (0.0012-0.0055%) in rice on the Swedish market [11]. There is still very little literature that discusses the usefulness of Rb in plants.

The types of essential heavy metals found in this study are Zn, Cu, Fe, and Mn, in certain quantities that are needed by living things, but if found in excessive amounts it will cause toxic effects. However, the mineral content found in this study showed no potential for causing poisoning. According to [12] and [3] types of heavy metals that are toxic to humans such as: arsenic (As), cadmium (Cd), copper (Cu), lead (Pb), mercury (Hg), nickel (Ni), (Zn), however, in the results this study found no type of heavy metal. From the description above, it can be assumed that the origin of the location and the environment of planting the sampled rice is still relatively safe and the rice has a reasonable mineral content for consumption.

#### IV. CONCLUSION

The mineral content for various types of rice that have been tested is varied. The highest mineral content was found in Kewal Bulu Hideung rice (3.884%), and the lowest was Sokan (1.677%). The mineral content found in succession from the largest to the smallest found in all rice tested was P, K, S, Ca, and Zn. The mineral content found only in certain varieties of rice is Mn, Fe, Rb, Si, Cu.

#### ACKNOWLEDGMENT

Thank you for the Islamic Development Bank which has funded this research as part of the RC Biotechnology participation in University of Sultan Ageng Tirtayasa, 2018. Thank you also said to the Shimadzu Laboratory, Osaka-Japan which facilitated this research.

#### REFERENCES

- [1] Z. Shahzad, H. Rouached, A. Rakha, "Combating mineral malnutrition through iron and zinc biofortification of cereals", *Compr. Rev. Food Sci. Food Saf.* 13:329-346, 2014

- [2] W.R. Rohaeni, E. Supriadi, U. Susanto, and T.D. Rosahdi, "Kandungan Fe dan Zn pada beras pecah kulit dan beras sosoh dari galur-galur padi toleran wereng batang cokelat (Fe and Zn content of brown rice and milled rice on brown planthopper tolerant rice lines)", *Jurnal Ilmu Pertanian Indonesia*, 21 (3): 172–176, 2016
- [3] D. Sundari, M. Hananto, dan Suharjo, "Kandungan logam berat dalam bahan pangan di kawasan industri kilang minyak, Dumai (Heavy metal in food ingredients in oil refinery industrial area, Dumai)", *Buletin Penelitian Sistem Kesehatan*, 19 (1): 55–61, 2016
- [4] T. Agustina, "Kontaminasi logam berat pada makanan dan dampaknya pada kesehatan", *Teknoboga*, 1(1): 53-65, 2014
- [5] M. Astawan, "Bahaya logam berat dalam makanan". Available on: <http://edukasi.kompas.com/read/2008/09/21/11254074/Bahaya.Logam.Berat.alam.Makanan>, 2008
- [6] S.D. Indrasari, P. Wibowo, A.A. Darajat, "Kandungan mineral beras varietas unggul baru" *Prossiding seminar nasional padi*. Source: [http://www.litbang.deptan.go.id/special/padi/bbpadi\\_2008\\_proeb412.pdf](http://www.litbang.deptan.go.id/special/padi/bbpadi_2008_proeb412.pdf), 2008
- [7] A.O. Oko, and S.I. Ugwu, "The proximate and mineral compositions of five major rice varieties in Abakaliki, South-Eastern Nigeria", *International Journal of Plant Physiology and Biochemistry*. 3(2): 25-27. <http://www.academicjournals.org/ijppb> ISSN-2141-2162, 2011
- [8] T. Iwai, M. Takahashi, O. Oda, Y. Terada., K.T. Yoshida, "Dynamic changes in the distribution of minerals in relation to phytic acid accumulation during rice seed development", *Plant Physiology Preview*. DOI: <https://doi.org/10.1104/pp.112.206573>, 2012
- [9] D.K. Verma, and P.P. Srivastav, "Proximate Composition, Mineral Content and Fatty Acids Analyses of Aromatic and Non-Aromatic Indian Rice" *Rice Science* 24(1):21-31· January 2017
- [10] T.H. Ansari, K. Iwasaki, T. Yoshida, Y. Yamamoto, and A. Miyazaki, "Status of nutrient elements in rice grain in relation to silicon accumulation pattern during grain filling", *Bangladesh Agron. J.* 19(2): 125-137 S, 2016
- [11] L. Jorhem, C. Åstrand, B. Sundström, M. Baxter, P. Stokes, J. Lewis, K.P. Grawé, "Elements in rice on the Swedish market: Part 2. Chromium, copper, iron, manganese, platinum, rubidium, selenium and zinc" *Food Addit. Contam.* 25:841-850, 2016
- [12] WHO, "WHO guidelines for indoor air: Selected pollutants: WHO Regional Office for Europe. Denmark, 2010

# Mineral Contents of Several Indonesian Rice Varieties

---

## ORIGINALITY REPORT

---

9%

SIMILARITY INDEX

5%

INTERNET SOURCES

6%

PUBLICATIONS

2%

STUDENT PAPERS

---

## MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

---

1%

★ [www.siftdesk.org](http://www.siftdesk.org)

Internet Source

---

Exclude quotes  On

Exclude matches  Off

Exclude bibliography  On

# Mineral Contents of Several Indonesian Rice Varieties

---

GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---