

Optimization of Ultrasonic— Assisted Extraction (UAE) Method Using Natural Deep Eutectic Solvent (NADES) to Increase Curcuminoid Yield from *Curcuma longa* L., *Curcuma xanthorrhiza*, and *Curcuma mangga* Va

Submission date: 08-May-2023 12:17PM (UTC+0700)
by Eka Sari

Submission ID: 2087220593


File name: molecules-27-06080_desi_s3_ui.pdf (1.48M)

Word count: 6099

Character count: 41329

Article

Optimization of Ultrasonic—Assisted Extraction (UAE) Method Using Natural Deep Eutectic Solvent (NADES) to Increase Curcuminoid Yield from *Curcuma longa* L., *Curcuma xanthorrhiza*, and *Curcuma mangga* Val.

Desy Rosarina ¹, Dimas Rafi Narawangsa ¹, Nabila Shaffa Rizky Chandra ¹, Eka Sari ^{2,3} and Heri Hermansyah ^{1,*} 

22

¹ Department of Chemical Engineering, Faculty of Engineering, Universitas Indonesia, Depok 16424, Indonesia

² Bioengineering & Biomedical Engineering, Research Centre CoE, Engineering Faculty, Sultan Ageng Tirtayasa University, Cilegon 42434, Indonesia

³ Chemical Engineering, Engineering Faculty, Sultan Ageng Tirtayasa University, Cilegon 42434, Indonesia

* Correspondence: heri.hermansyah@ui.ac.id

16

Abstract: This study aims to optimize ultrasonic-assisted natural deep eutectic solvents (NADES) based extraction from *C. longa*. Choline chloride-lactic acid (CCLA-H₂O = 1:1, b/v) was used to investigate the impact of various process parameters such as solvent's water content, solid loading, temperature, and extraction time. The optimal yield of 79.635 mg/g of *C. longa* was achieved from extraction in 20% water content NADES with a 4% solid loading in 35 °C temperature for 1 h. Peleg's model was used to describe the kinetics of the optimized ultrasonic-assisted extraction (UAE) method, and the results were found to be compatible with experimental data. The optimum conditions obtained from *C. longa* extraction were then used for the extraction of *C. xanthorrhiza* and *C. mangga*, which give yields of 2.056 and 31.322 mg/g, respectively. Furthermore, n-hexane was utilized as an anti-solvent in the separation process of curcuminoids extract from *C. longa*, *C. xanthorrhiza*, and *C. mangga*, which gave curcuminoid recovery of 39%, 0.74%, and 27%, respectively. Solidification of curcuminoids was also carried out using the crystallization method with n-hexane and isopropanol. However, the solution of CCLA and curcuminoids formed a homogeneous mixture with isopropanol. Hence, the curcuminoids could not be solidified due to the presence of NADES in the extract solution.

46

Keywords: curcumin; turmeric; extraction; ultrasonic-assisted extraction; natural deep eutectic solvents



Citation: Rosarina, D.; Narawangsa, D.R.; Chandra, N.S.R.; Sari, E.; Hermansyah, H. Optimization of Ultrasonic—Assisted Extraction (UAE) Method Using Natural Deep Eutectic Solvent (NADES) to Increase Curcuminoid Yield from *Curcuma longa* L., *Curcuma xanthorrhiza*, and *Curcuma mangga* Val. *Molecules* **2022**, *27*, 6080. <https://doi.org/10.3390/molecules27186080>

Academic Editor: Giancarlo Cravotto

Received: 14 August 2022

Accepted: 15 September 2022

Published: 18 September 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Curcuminoids are hydrophobic polyphenols found in rhizomes of plants from the *Zingiberaceae* family that are widely cultivated in Asia, including India and Indonesia. Curcuminoids are the main pigment of the plants, giving their rhizomes a bright, yellow-orange color. These compounds consist of three forms of polyphenols, namely, curcumin (C₂₁H₂₀O₆), demethoxycurcumin (C₂₀H₁₈O₅), and bisdemethoxycurcumin (C₁₉H₁₆O₄). The highest curcuminoid content was found in the rhizomes of the *C. longa* (3–8%), *C. xanthorrhiza* (2%), and *C. mangga* (3%) [1–3].

Curcuminoids are known for having many medical benefits. The pharmacological effects of curcuminoids, such as antiinflammation, antioxidant, antibacterial, anticancer, and anti-allergy properties, have been medically proven, both *in vitro* and *in vivo* [4]. Curcumin is also known to inhibit angiogenesis and tumor development and to induce apoptosis [5,6]. In addition, based on *in silico* studies, curcumin is also known to have high affinity for the spike (s) glycoprotein of SARS-CoV-2 and ACE2 which indicates its potential antiviral activity in preventing the transmission of COVID-19 [7].

According to the Joint FAO/WHO Expert Committee, naturally extracted curcuminoids are preferred over chemically synthesized curcuminoids as a food additive. This has

led to numerous studies on the most suitable extraction methods for this compound [8–12]. Some conventional methods such as maceration, Soxhlet, and reflux can be used to extract curcumin successfully. However, those methods have some major disadvantages: long extraction time, the use of large amounts of organic solvents, and high temperature that can cause the degradation of curcuminoids, leading to poor efficiency of extraction and high energy consumption.

In recent years, modern methods such as ultrasonic-assisted extraction (UAE) have been researched for a faster and more cost-effective extraction of natural resources [10,12]. This method utilizes ultrasonic waves ranging from 20 kHz to 2000 kHz to generate shock waves and various physical phenomena causing the cavitation, i.e., the formation, growth, and explosion, of the cavitation bubbles. The cavitation process results in intense local heating and pressure. Based on the hotspots mechanism, the bubble burst increases at a local temperature of more than 5273 K and a pressure of about 2000 atm [13]. However, this phenomenon does not cause heat damage to the phytochemical because the bubble burst occurs at a fast time with cooling rates in the range of 10^{10} K/s [13]. These extreme conditions break chemical bonds, thus helping the diffusion of solvents to plant cell walls. Therefore, the extraction can be carried out faster even though the process occurs at room temperature [14].

For the extraction of curcuminoids, organic solvents are commonly used. Curcuminoids can easily dissolve in organic solvents such as ethanol, methanol, ethyl acetate, and acetone [8]. However, although the organic solvents have important characteristics for the extraction and the dissolution of natural resources, they have limitations, especially when used in extensive amounts, such as toxicity, non-biodegradability, environmental pollution, lack of selectivity for the desired compound, flammability, and leaving hazardous residue in the extract [15]. Therefore, developing green and sustainable solvents as an alternative to organic solvents is expected to solve these problems.

NADES (natural deep eutectic solvent) is a green solvent composed of natural primary metabolites such as sugars, alcohols, amino acids, amines, and carboxylic acids [16]. NADES is formed by mixing quaternary ammonium salts, i.e., hydrogen bond acceptor (HBA), which forms hydrogen bonds with hydrogen bond donors (HBD) [17]. This solvent is called eutectic because a eutectic point will be formed when the two components are mixed in the correct ratio [18]. Compared to organic solvents, NADES has advantages, such as biocompatibility, biodegradability, low toxicity, and low-cost [15]. However, NADES has a high viscosity, which results in its slow diffusivity in the plant matrix during extraction. Therefore, adding water to the mixture of HBA and HBD plays an important role in NADES-based extraction.

In this research, the UAE method is optimized by applying four variations: water content in solvent (NADES), solid loading (w/v %), extraction temperature, and time. These variations are carried out based on the one-variable-at-a-time (OVAT) principle, meaning that when the first parameter is varied, the other parameters are set constant. A comparison study has also been carried out between the Soxhlet and UAE-optimized methods. In addition, a kinetics study was conducted using Peleg's model to evaluate the extraction kinetics parameters and their compatibility with experimental data for UAE extraction of curcuminoids from *C. longa*. The research then continued to optimize the yield of curcuminoid extract from *C. xanthorrhiza* and *C. mangga* based on the previously optimized parameter obtained. The separation and crystallization methods were also used to produce high purity curcuminoids and maintain the curcuminoid's physical and chemical stability. Curcuminoids extract is quantified with the absorbance test using UV-VIS spectrophotometry.

2. Results and Discussion

2.1. Soxhlet Method

The Soxhlet method resulted in 88.476 mg curcuminoid/g from *C. longa* (8.8% w/w). The obtained yield was considered as reference, i.e., 100%. Note that the curcuminoid

obtained from this experiment is a crude extract, since it is not pure and might still contain some other polyphenols and many other minor compounds. Viscous extract obtained from this process is 1.56 g (31.2% *w/w*) from the original 5 g *C. longa* powder.

2.2. Optimized UAE Method

2.2.1. Water Content (%) Variation

The major drawback of using NADES as an extraction solvent is its high viscosity which can inhibit the diffusion of the solvent to the cell matrix [12]. The dissolution of NADES with distilled water aims to reduce the high viscosity of NADES [15]. Therefore, in this study, the effect of adding water into NADES (20%, 25%, and 30%) was investigated.

The best variation obtained is 20% water content in NADES at 60 min with 43.83 mg/g curcuminoid extract yield. The other two variations, 25% and 30%, gave lower results at 60 min extraction time: 32.58 and 30.74 mg/g, respectively. The complete extraction data will be shown in Figure 1.




Figure 1. Curcumin extract yield to time graph for water content variation.

As shown in Figure 1, the 20% water content in NADES gave the highest yield of curcuminoids. In ultrasonic extraction, viscosity and surface tension play a significant role [10]. The decrease in viscosity and surface tension causes the formation of cavitation explosions more easily. Fluids with lower viscosity tend to form stable bursts of bubbles and cavitation with a larger magnitude than those with high viscosity [19]. Thus, this helps facilitate easier diffusion of the solvent into the plant sample. However, adding water content of as much as 5% successively into NADES causes a decrease in the yield of curcuminoids. Gabriele (2019), who also investigated the effect of adding water content to NADES, stated that the hydrogen bond between HBA (hydrogen bond acceptor) and HBD (hydrogen bond donor) would weaken with the addition of water content up to 50% [20]. 75% water content, the bond is completely broken. Therefore, in this experiment, 20% water content in NADES is the optimal condition.

2.2.2. Solid Loading (%) Variation

Mass transfer at the solid–liquid interface is influenced by the proportion of solids and the amount of solvent under specific extraction operating conditions. Therefore, in this study, the effect of % (*w/v*) solid loading (7%, 5%, and 4%) with extraction operating conditions of 20% water content in NADES, temperature 35 °C, particle size 0.25 mm, power 60% (6 s on, 4 s off), for 60 min was studied.

As shown in Figure 2, the best variation obtained is 4% solid loading at 60 min with 79.67 mg/g curcuminoid extract yield. The other two variations, 5% and 7%, gave lower results at 60 min extraction time: 64.55 and 43.83 mg/g, respectively.




Figure 2. Curcumin extract yield to time graph for solid loading variation.

The yield of curcuminoid extract increased alongside the decrease of solid loading. At a fixed amount of solid matrix, the more solvent volume used, the greater the concentration gradient between the plant cell matrix and the extraction medium. This increases the driving force to release curcuminoids from the sample matrix to the solvent. In addition, a decrease in the solids loading in the solvent also causes cavitation at a lower cavitation threshold [21]. This would significantly increase the extraction rate.

2.2.3. Temperature Variation

Based on Figure 3, the curcuminoids yield obtained increased alongside the increase in temperature from 35 °C to 55 °C. After 60 min of sonication, the maximum curcuminoids yield of 79.64, 80.64, and 84.27 mg/g was achieved from extraction temperatures of 35 °C, 45 °C, and 55 °C, respectively.




Figure 3. Curcumin extract yield to time graph for temperature variation.

The yield of curcuminoid extract increases alongside the increase in temperature. Higher temperature increases the extraction rate because it gives the active compound more kinetic energy. According to diffusivity and solubility of solute, the temperature increase would increase both parameters and thus increase extraction effectivity [22]. Higher temperature would also increase solid material's pore size, increasing diffusivity further [10]. However, high temperatures can lead to curcumin degradation in the material. This happened at 55 °C temperature, at which the mixture color changed from bright yellow-orangish to dark brown. The dark brown color indicated other compounds, such as ferulic acid, vanillic acid, and vanillin [23,24]. Because of this, 55 °C temperature cannot

be used as the optimal condition. Instead, the 35 °C temperature would be chosen since it gave a slightly lower result than the 45 °C and 55 °C, and it used less energy than both temperatures.

2.2.4. Effect of Extraction Time

The extraction process depends on reaction time, and in this study, the extraction process was also observed with variation in extraction time. The longer the extraction was carried out, the more yield was obtained. As shown in Figures 1–3, the extraction rate was very fast in the first 20 min. After that, the yields obtained increased gradually up to 1 h. The curcuminoid concentration gradient between the solvent and *C. longa* powder was extensive at the beginning of the extraction process. In addition, the extraction was facilitated by the ease of extraction from the outer matrix during the initial period. As time goes by, the concentration gradient is lower, and the extraction progress is slower because the remaining curcuminoids are in the core of the solid matrix, so this is influenced by diffusion from the solvent. Therefore, there is a limit on the curcuminoids that can be extracted. For every variation, an extraction capacity shows approximately how much curcuminoids can be extracted. The yield will not surpass the extraction capacity, no matter how long the extraction is carried out. The capacity can be calculated empirically using Peleg's model.

2.2.5. Optimized UAE and Soxhlet Method Comparison

The yield of curcuminoids obtained from the Soxhlet method is considered a value of 100%, which is 88.48 mg/g. In the extraction using the UAE method, a yield of 79.64 mg/g (90%) was obtained with the extraction process only at ± 35 °C for 1 h.

Figure 4 shows how the productivities obtained from extraction with Soxhlet and UAE compare. Although Soxhlet obtained a higher yield, it required a longer time, more solvent, and energy compared to the optimized UA method. Therefore, the productivity of the UAE method (79.64 mg/g·h) is better than that of Soxhlet (7.37 mg/g·h).




Figure 4. Comparison of extraction productivity obtained from Soxhlet and UAE.

The shear force generated due to cavitation along with the shock wave induces physical damage to the plant cell wall, thereby facilitating the release of the compound to be extracted [25]. Diffusion of the solvent also increases with swelling of the plant matrix and hydration by the solvent [26,27]. Ultrasonic wave also increases mass transfer, so solvent diffusion into the plant cell matrix is more effective [10]. Therefore, the UAE only takes a shorter time in the extraction process and requires less energy compared to Soxhlet.

2.3. Curcuminoid Extract from *C. xanthorrhiza* and *C. mangga*

The extraction of curcuminoids from *C. xanthorrhiza* and *C. mangga* with UAE was carried out using the optimized parameters previously obtained from *C. longa* extraction. Based on the experiment, the UAE-optimized method gave yields of 2.06 and 31.32 mg/g for *C. xanthorrhiza* and *C. mangga*, respectively. The comparison is shown on Figure 5.




Figure 5. Curcumin extract yield comparison.

The yields obtained from *C. xanthorrhiza* and *C. mangga* were significantly lower than the one obtained from *C. longa*. *C. xanthorrhiza* has a curcuminoid content up to 1–2% of its total weight [2], while *C. mangga* has up to 3% [3]. *C. longa* has around 3–8% [1], which makes it the highest source of curcuminoid compared to the other two. *C. xanthorrhiza* does not contain bisdemethoxycurcumin [28], which is easier to extract compared to curcumin and demethoxycurcumin [29]. This makes *C. longa*, which contains all three compounds, give the highest yield. Therefore, *C. longa* is the best option to obtain curcuminoid extract.

2.4. Separation and Crystallization Methods

In optimal conditions, separation and crystallization methods were carried out on the curcuminoid extract obtained from *C. longa*, Javanese turmeric, and *C. mangga*. In this study, curcuminoids separated from oleoresin using the anti-solvent precipitation method, where n-hexane was used as antisolvent as curcuminoids are insoluble in it. This process gives curcuminoids recoveries of 39%, 0.74%, and 27% from *C. longa*, *C. xanthorrhiza*, and *C. mangga*, respectively.

The crystallization process is carried out after the separation process to obtain the curcuminoids in solid form. The mixture of isopropanol-hexane (1:1.5) at 40 °C was used in the solidification process. This process gave curcuminoids recoveries of 22%, 1.29%, and 13% from *C. longa*, *C. xanthorrhiza*, and *C. mangga*, respectively. However, curcuminoids extract cannot be solidified directly from NADES solution using isopropanol-hexane.

The result showed that both methods can recover some curcuminoids in the extract, which means both methods increase the concentration of curcuminoid extracted. The separation method resulted in a bigger recovery for *C. longa*, but the crystallization method had lower result. The opposite happened to *C. xanthorrhiza* and *C. mangga*, where the curcuminoid recovery obtained from the crystallization method was bigger than that of the separation method.

The recovery of curcuminoid in the extract can be explained by the antisolvent used in the process. Curcuminoid extract was dissolved in n-hexane which acted as antisolvent. Difference of polarity caused the curcuminoid to precipitate at the bottom of the mixture while being separated from other impurities such as oleoresin [11].

The crystallization process then was attempted to solidify the extract and gain pure curcuminoid powder, but it was unsuccessful. The mixture between NADES and the

antisolvent–solvent solution (n-hexane and isopropanol) caused a competition in which both solutions tried to dissolve the curcuminoid. This resulted in precipitation of curcuminoid extract, in which no solids were formed, but some curcuminoids could still be further recovered [11].

2.5. Kinetic Model

In this study, the extraction process is modeled with a semi-empirical kinetic model named Peleg's model. The simplified equation is shown below [12].

$$M_t = M_0 + \frac{t}{K_1 + K_2 t} \quad (1)$$

where:

M_t —final concentration of curcumin extract at time t ($\text{mg}_{\text{curcuminoid}} / \text{g}_{\text{material}}$);

M_0 —initial concentration of curcumin extract ($\text{mg}_{\text{curcuminoid}} / \text{g}_{\text{material}}$);

K_1 —Peleg's rate constant ($\text{min g}_{\text{material}} / \text{mg}_{\text{curcuminoid}}$);

K_2 —Peleg's capacity constant ($\text{g}_{\text{material}} / \text{mg}_{\text{curcuminoid}}$);

t —extraction time (min).

The equation is made into linear regression equation, with $1/M_t$ as y , $1/t$ as x , K_1 as gradient, and K_2 as intercept. The equation is shown below.

$$\frac{1}{M_t} = K_1 \frac{1}{t} + K_2 \quad (2)$$

The obtained data would be plotted on a graph to obtain the linear regression equation and coefficient of determination for the models. The model has some kinetic parameters, such as K_1 (Peleg's rate constant), K_2 (Peleg's capacity constant), B_0 (initial extraction rate), and C_e (curcuminoids concentration at equilibrium) [30].

Peleg's model was used to describe the extraction kinetic of the optimized UAE method. *C. longa* extraction results are used to make this model. Each variation would have one Peleg's model to determine their own parameters. The Peleg's models created from three variations are shown in Figures 6–8.




Figure 6. Peleg's model for water content variation.




Figure 7. Peleg's model for solid loading variation.




Figure 8. Peleg's model for temperature variation.

Compatibility of kinetics study with experiment data was determined using RMSE, adjusted R^2 , and E (%) value. Table 1 revealed the results obtained for Peleg's model.

Table 1. RMSE, R^2 , and E% value for Peleg's model.

Variables		RMSE	R^2	E (%)
Water content (%)	20	0.43	0.97	0.05%
	25	0.26	0.96	0.01%
	30	0.32	0.96	0.01%
Solid loading (%)	7	0.43	0.97	0.05%
	5	0.44	0.99	0.05%
	4	0.35	0.99	0.02%
Extraction temperature (°C)	35	0.35	0.99	0.02%
	45	0.42	0.99	0.04%
	55	0.38	0.99	0.05%
Average		0.38	0.97 *	0.03

* Adjusted R^2 .

Table 2 shows the kinetic parameters for Peleg's model, which include K_1 (Peleg's rate constant), K_2 (Peleg's capacity constant), B_0 (initial extraction rate), and C_e (curcuminoids concentration at equilibrium).

Table 2. Kinetics parameter for Peleg's model.

Variables		K_1 (g min/mg)	K_2 (g/mg)	B_0 (mg/g min)	C_e (mg/g)
Water content (%)	20	0.19	0.02	5.15	49.26
	25	0.13	0.03	7.70	35.21
	30	0.15	0.03	6.67	31.65
Solid loading (%)	7	0.19	0.02	5.15	49.26
	5	0.27	0.01	3.76	86.96
	4	0.22	0.01	4.50	109.89
Extraction temperature (°C)	35	0.22	0.01	4.50	109.89
	45	0.20	0.01	5.02	107.53
	55	0.18	0.01	5.73	114.94

The Peleg's model showed a relation between extraction time and the extract concentration in solvent. Figures 6–8 showed an increase of curcuminoid extract yield alongside the increase of extraction time. The errors in these models are very miniscule and can be ignored. The model showed a very good coefficient of determination (R^2), which means a good correlation between each data. The parameters shown can illustrate how well the extraction of each variations went. The bigger the C_e , more curcuminoid extract can be obtained from the variation that is being carried out. Since this model is linear, it showed that the 55 °C variation has the highest C_e . However, this is not the case since the 55 °C variation has a risk of curcumin degradation because of high temperature, indicated by color change of the mixture from yellow-orangish to dark brown. While the model showed a good correlation with the experimental data, it cannot show how the extraction went in the experiment. That is why the optimal variation is the 35 °C temperature.

3. Materials and Methods

3.1. Materials

Fresh *C. longa* was purchased at Tangerang (Banten, Indonesia) local market. Fresh *C. mangga* was commercially purchased from a local business at Garut (West Java, Indonesia). Fresh *C. xanthorrhiza* was purchased at local market in Jakarta Selatan (DKI Jakarta, Indonesia).

Pure curcuminoids standard was commercially purchased from a retailer at Bantul (Yogyakarta, Indonesia). Choline chloride (95%), lactic acid (90%), analytical grade ethanol (>99%), hexane (>98%), and isopropanol (>99.8%) were obtained from Merck (Darmstadt, Germany). A nylon membrane filter with a pore size of 0.45 μm was obtained from Merck Millipore (Carrigtwohill, Ireland). Whatman filter paper (No. 42) with 90 mm pore size was obtained from Whatman (Dassel, Germany).

3.2. Preparation of NADES

In this study, NADES was prepared with the heating and stirring method. Choline chloride (HBA) and lactic acid (HBD) were mixed with a 1:2 molar ratio. Water content in NADES was adjusted by adding aquadest. The components were mixed in a beaker glass at 70 °C with constant stirring until clear, homogenous liquid formed. NADES was then stored in a sealed glass at room temperature until use.

3.3. Extraction Method

3.3.1. Soxhlet Extraction

Soxhlet extraction was utilized as the reference method to compare the optimization results performed on the ultrasound-assisted extraction (UAE) method. The procedure for

this method was carried out based on Patil et al. (2020) [12]. A total of 5 g of *C. longa* powder in the thimble was extracted using ethanol (96%) with a ratio of 1:50 (*w/v*). The extraction process was carried out for 12 h at temperature of ± 70 °C. The curcuminoid yield obtained was a reference result (100%).

3.3.2. Ultrasonic-Assisted Extraction

Ultrasound-Assisted Extraction was carried out with a NADES-based solvent, choline chloride: lactic acid (1:2), with 250 W rated output power and a frequency of 22 kHz. The extraction was performed with a sonicator probe operated at pulsed mode with a 60% duty cycle (6 s on and 4 s off). The ultrasonic waves were transmitted directly to the mixture for 1 h, with data collection every 10 min. The samples then were diluted, centrifuged, and analyzed quantitatively using UV-VIS spectrophotometry. The effect of water content (%) in NADES, solid loading, extraction temperature, and time were investigated to optimize the process.

3.4. Curcumin Quantification

Spectrophotometry UV-VIS was used to analyze the curcuminoids concentration from the samples. The measurements were carried out at a wavelength of 415 nm, giving the absorbance data. The concentrations of curcuminoids (ppm) were calculated using a standard curve of 95% pure curcuminoids and consequently calculated as the yield of curcuminoids (mg/g).

3.5. Curcumin Extract Separation

In the present work, curcuminoids and oleoresin were separated using an antisolvent technique with n-hexane. The viscous extract obtained from the optimized UAE method was centrifuged at 6000 rpm for 50 min, and the supernatant was stored in another tube. A known volume of supernatant was mixed with hexane (1:25, *v/v*) and allowed to stand for 24 h. After 24 h, the solution was stirred with a magnetic stirrer at 600 rpm for 3 h. The precipitated mixture was separated from oleoresin by centrifuging the solution at 1000 rpm for 10 min. The recovery of curcuminoids was analyzed quantitatively using spectrophotometry UV-VIS at 415 nm wavelength.

3.6. Curcumin Extract Crystallization

The separated curcuminoids from a previous process were further purified employing the crystallization method used by Pawar et al. (2018) [8]. Curcuminoid extract was added to 10 mL of isopropanol and n-hexane mixture (4:6, *v/v*) at 40 °C. The mixture and extract were then cooled in a freezer (temperature below 10 °C) for approximately 1 h. The sediment then was separated from the filtrate using a vacuum filter and then analyzed using UV-VIS spectrophotometry.

4. Conclusions

The optimal condition for NADES based UAE in this research is 20% water content, 4% solid loading, 35 °C temperature, and 60 min extraction time. This condition resulted in 79.636 mg curcuminoid extract/g *C. longa*. Peleg's model was used to describe the kinetics of the optimized UAE method, and the results were found to be compatible with experimental data. Based on the yield obtained, the use of solvents, temperature, and extraction time, the optimized UAE method can be chosen as an alternative to the Soxhlet method. Furthermore, *C. longa* is the best material from which to obtain curcumin extract, since *C. xanthorrhiza* and *C. mangga* gave a significantly lower result. The separation and crystallization methods are worth carrying out with *C. longa* extract since this extract gave the highest recoveries of curcuminoid, which were 39% from separation and 22% from crystallization. However, curcuminoids could not be solidified due to the presence of NADES in the extract solution.

Author Contributions: H.H.: conceptualization, funding acquisition, project administration, resources, supervision; D.R.N.: conceptualization, project administration, formal analysis, investigation, methodology, visualization, writing—original draft, writing—review and editing; N.S.R.C.: conceptualization, project administration, formal analysis, investigation, methodology, visualization, writing—original draft, writing—review and editing; D.R.: validation; E.S.: validation. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by Kementerian Pendidikan, Kebudayaan, Riset, dan Teknologi Republik Indonesia (NKB-1057/UN2.RST/HKP.05.00/2022).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: The authors would like to express gratitude to Bioprocess Lab in Department of Chemical Engineering for in-kind support, and to Direktorat Riset, Teknologi, dan Pengabdian Kepada Masyarakat, Direktorat Jenderal Pendidikan Tinggi, Riset, dan Teknologi, and Kementerian Pendidikan, Kebudayaan, Riset, dan Teknologi Republik Indonesia for funding this research.

Conflicts of Interest: The authors declare no conflict of interest.

Sample Availability: Samples of the curcumin extracts are not available from the authors.

References

1. Goel, A.; Kunnumakkara, A.B.; Aggarwal, B.B. Curcumin as "Curecumin": From Kitchen to Clinic. *Biochem. Pharmacol.* **2008**, *75*, 787–809. [[CrossRef](#)] [[PubMed](#)]
2. Basalmah, R.S. Optimalisasi Kondisi Ekstraksi Kurkuminoid Temulawak: Waktu, Suhu Dan Nisbah. Bachelor's Thesis, Institut Pertanian Bogor, Bogor, Indonesia, 2006.
3. Jantan, I.B.; Ahmad, A.S.; Ali, N.A.M.; Ahmad, A.R.; Ibrahim, H. Chemical Composition of the Rhizome Oils of Four Curcuma Species from Malaysia. *J. Essent. Oil Res.* **1999**, *11*, 719–723. [[CrossRef](#)]
4. Dei Cas, M.; Ghidoni, R. Dietary Curcumin: Correlation between Bioavailability and Health Potential. *Nutrients* **2019**, *11*, 2147. [[CrossRef](#)] [[PubMed](#)]
5. Choundhuri, T.; Pal, S.; Das, T.; Sa, G. Curcumin selectivity induces apoptosis in deregulated cyclin D1-expressed cells at G2 phase of cell cycle in a p53-99 dependent manner. *J. Biol. Chem.* **2005**, *280*, 20059–20068. [[CrossRef](#)]
6. Lin, Y.G.; Kunnumakkara, A.B.; Nair, A.; Merritt, W.M.; Han, L.Y.; Armaiz-Pena, G.N.; Kamat, A.A.; Spannuth, W.A.; Gershenson, D.M.; Lutgendorf, S.K.; et al. Curcumin inhibits tumor growth and angiogenesis in ovarian carcinoma by targeting the nuclear factor κ B pathway. *Clin. Cancer Res.* **2007**, *13*, 3423–3430. [[CrossRef](#)]
7. Patel, A.; Rajendran, M.; Shah, A.; Patel, H.; Pakala, S.B.; Karyala, P. Virtual screening of curcumin and its analogs against the spike surface glycoprotein of SARS-CoV-2 and SARS-CoV. *J. Biomol. Struct. Dyn.* **2022**, *40*, 5138–5146. [[CrossRef](#)]
8. Pawar, H.A.; Gavasane, A.J.; Choudhary, P.D. A Novel and Simple Approach for Extraction and Isolation of Curcuminoids from Turmeric Rhizomes. *Nat. Prod. Chem. Res.* **2018**, *6*, 1–4. [[CrossRef](#)]
9. Jiang, T.; Ghosh, R.; Charcosset, C. Extraction, Purification and Applications of Curcumin from Plant Materials-A Comprehensive Review. *Trends Food Sci. Technol.* **2021**, *112*, 419–430. [[CrossRef](#)]
10. Shirsath, S.R.; Sable, S.S.; Sonawane, S.H.; Saini, D.R.; Gogate, P.R. Intensification of Extraction of Curcumin from *Curcuma amada* Using Ultrasound Assisted Approach: Effect of Different Operating Parameters. *Ultrason. Sonochemistry* **2017**, *38*, 437–445. [[CrossRef](#)]
11. Rachmaniah, O.; Muhsin, M.; Widya, A. and Rachimoellah, M. Process Development for The Enrichment of Curcuminoids in The Extract of Ionic Type Of NADES. *IOP Conf. Ser. Mater. Sci. Eng.* **2020**, *732*, 012005. [[CrossRef](#)]
12. Patil, S.S.; Pathak, A.; Rathod, V.K. Optimization and Kinetic Study of Ultrasound Assisted Deep Eutectic Solvent Based Extraction: A Greener Route for Extraction of Curcuminoids from *Curcuma longa*. *Ultrason.-Sonochemistry* **2021**, *70*, 1–10. [[CrossRef](#)] [[PubMed](#)]
13. Ashiri, R. Obtaining a novel crystalline/amorphous core/shell structure in barium titanate nanocrystals by an innovative one-step approach. *R Soc. Chem.* **2015**, *5*, 48281–48289. [[CrossRef](#)]
14. Handa, S.; Khanuja, S.P.S.; Longo, G.; Rakesh, D.D. *Extraction Technologies for Medicinal and Aromatic Plants*; ICS UNIDO: Trieste, Italy, 2008.
15. Dai, Y.; Witkamp, G.J.; Verpoorte, R.; Choi, Y.H. Natural deep eutectic solvents as a new extraction media for phenolic metabolites in *Carthamus tinctorius* L. *Anal. Chemi.* **2013**, *13*, 6272–6278. [[CrossRef](#)] [[PubMed](#)]
16. Ruesgas-Ramon, M.; Figueroa-Espinoza, M.C.; Durand, E. Application of deep eutectic solvents (DES) for phenolic compounds extraction: Overview, challenges, and opportunities. *J. Agric. Food Chem.* **2017**, *65*, 3591–3601. [[CrossRef](#)]

17. Durand, E.; Lecomte, J.; Villeneuve, P. Deep eutectic solvents: Synthesis, application, and focus on lipase-catalyzed reactions. *Eur. J. Lipid Sci. Technol.* **2013**, *115*, 379–385. [[CrossRef](#)]
18. Ali, E.; Kali, M.K.H.; Mulyono, S.; Alnashef, I.; Fakeeha, A.; Mjalli, F.; Hayyan, A. Solubility of CO₂ in deep eutectic solvents: Experiments and modelling using the Peng–Robinson equation of state. *Chem. Eng. Res. Des.* **2014**, *94*, 1898–1906. [[CrossRef](#)]
19. Helfield, B.; Black, J.J.; Qin, B.; Pacella, J.; Chen, X.; Villanueva, F.S. Fluid Viscosity Affects the Fragmentation and Inertial Cavitation Threshold of Lipid-Encapsulated Microbubbles. *Ultrasound Med. Biol.* **2016**, *42*, 782–794. [[CrossRef](#)]
20. Gabriele, F.; Chiarini, M.; Germani, R.; Tiecco, M.; Spreti, N. Effect of water addition on choline chloride/glycol deep eutectic solvents: Characterization of their structural and physicochemical properties. *J. Mol. Liq.* **2019**, *291*, 111301. [[CrossRef](#)]
21. Santos, H.M.; Lodeiro, C.; Capelo-Martnez, J. The Power of Ultrasound, Ultrasound in Chemistry. *Ultrasound Chem. Anal. Appl.* **2009**, *1*, 1–16. [[CrossRef](#)]
22. Yue, L.; Zhang, F.; Wang, Z. Ultrasound-Assisted Extraction of Capsaicin from Red Peppers and Mathematical Modelling. *Sep. Sci. Technol.* **2012**, *47*, 124–130. [[CrossRef](#)]
23. Suresh, D.; Gurudutt, K.N.; Srinivasan, K. Degradation of Bioactive Spice Compound: Curcumin During Domestic Cooking. *Eur. Food Res. Technol.* **2009**, *228*, 807–812. [[CrossRef](#)]
24. Tensiska; Nurhadi, B.; Isfron, A.F. Kestabilan Warna Kurkumin Terenkapsulasi dari Kunyit (*Curcuma Domestica* Val.) Dalam Minuman Ringan dan Jelly pada Berbagai Kondisi Penyimpanan. *Bionatura-J. Ilmu-Ilmu Hayati Dan Fis.* **2012**, *14*, 198–206. Available online: <http://jurnal.unpad.ac.id/bionatura/article/view/7461> (accessed on 20 April 2022).
25. Vinatoru, M. An overview of ultrasonically assisted extraction of bioactive principles from herbs. *Ultrason. Sonochemistry* **2001**, *8*, 303–313. [[CrossRef](#)]
26. Toma, M.; Vinatoru, M.; Paniwnyk, L.; Mason, J. Investigation of the effects of ultrasound on vegetal tissues during solvent extraction. *Ultrason. Sonochemistry* **2001**, *8*, 137–142. [[CrossRef](#)]
27. Jerman, T.; Trebse, P.; Vodopivec, B.M. Ultrasound-assisted solid liquid extraction (USLE) of olive fruit (*Olea europaea*) phenolic compounds. *Food Chem.* **2010**, *123*, 175–182. [[CrossRef](#)]
28. Sembiring, B.B.; Ma'mun, M.; Ginting, E.I. Pengaruh Kehalusan Bahan dan Lama Ekstraksi Terhadap Mutu Ekstrak Temulawak (*Curcuma xanthorrhiza*, Roxb). *Bul. Penelit. Tanam. Rempah Dan Obat* **2006**, *17*, 53–58. Available online: <http://repository.pertanian.go.id/handle/123456789/3805> (accessed on 20 April 2022).
29. Setyowati, A.; Suryani, C.L. Peningkatan Kadar Kurkuminoid Dan Aktivitas Antioksidan Minuman Instan Temulawak Dan Kunyit. *Agritech* **2013**, *33*, 363–370. [[CrossRef](#)]
30. Jokić, S.; Velić, D.; Bilić, M.; Bucić-Kojić, A.; Planinić, M.; Tomas, S. Modelling of the Process of Solid-Liquid Extraction of Total Polyphenols from Soybeans. *Czech J. Food Sci.* **2010**, *28*, 206–212. [[CrossRef](#)]

Optimization of Ultrasonic—Assisted Extraction (UAE) Method Using Natural Deep Eutectic Solvent (NADES) to Increase Curcuminoid Yield from *Curcuma longa* L., *Curcuma xanthorrhiza*, and *Curcuma mangga* Va

ORIGINALITY REPORT

16%

SIMILARITY INDEX

9%

INTERNET SOURCES

14%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Sujata S. Patil, Ajay Pathak, Virendra K. Rathod. "Optimization and kinetic study of ultrasound assisted deep eutectic solvent based extraction: A greener route for extraction of curcuminoids from *Curcuma longa*", *Ultrasonics Sonochemistry*, 2021
Publication 3%
- 2 Rui Wang, Kaiqing Zhang, Ivan V. Kozhevnikov. "Ultrasonic Auxiliary Ozone Oxidation-Extraction Desulfurization: A Highly Efficient and Stable Process for Ultra-Deep Desulfurization", *Molecules*, 2022
Publication 2%
- 3 S.R. Shirsath, S.S. Sable, S.G. Gaikwad, S.H. Sonawane, D.R. Saini, P.R. Gogate. "Intensification of extraction of curcumin from *Curcuma amada* using ultrasound assisted approach: Effect of different operating parameters", *Ultrasonics Sonochemistry*, 2017 2%

4	bjo.bmj.com Internet Source	1 %
5	bioresources.cnr.ncsu.edu Internet Source	<1 %
6	Madhulekha Rakshit, Prem P. Srivastav, Kanishka Bhunia. "Kinetic modeling of ultrasonic - assisted extraction of punicalagin from pomegranate peel", Journal of Food Process Engineering, 2020 Publication	<1 %
7	Sujata S. Patil, Virendra K. Rathod. "Synergistic Effect of Ultrasound and Three Phase Partitioning for the Extraction of Curcuminoids from Curcuma longa and its Bioactivity Profile", Process Biochemistry, 2020 Publication	<1 %
8	assets.ctfassets.net Internet Source	<1 %
9	www.coursehero.com Internet Source	<1 %
10	Adepu K. Kumar, Bhumika S. Parikh, Mohanty Pravakar. "Natural deep eutectic solvent mediated pretreatment of rice straw: bioanalytical characterization of lignin extract and enzymatic hydrolysis of pretreated	<1 %

biomass residue", Environmental Science and Pollution Research, 2015

Publication

11

Ramesh Sapkota, Paul Marshall. "Gas-Phase Chemistry of 1,1,2,3,3,4,4-Heptafluorobut-1-ene Initiated by Chlorine Atoms", Molecules, 2022

Publication

<1 %

12

deepblue.lib.umich.edu

Internet Source

<1 %

13

ikee.lib.auth.gr

Internet Source

<1 %

14

www.scientific-publications.net

Internet Source

<1 %

15

Andrés González de Castilla, Jan Philipp Bittner, Simon Müller, Sven Jakobtorweihen, Irina Smirnova. " Thermodynamic and Transport Properties Modeling of Deep Eutectic Solvents: A Review on g -Models, Equations of State, and Molecular Dynamics ", Journal of Chemical & Engineering Data, 2019

Publication

<1 %

16

Milena Ivanović, Peter Krajnc, Aleš Mlinarič, Maša Islamčević Razboršek. "Natural Deep Eutectic Solvent-Based Matrix Solid Phase Dispersion (MSPD) Extraction for Determination of Bioactive Compounds from

<1 %

Sandy Everlasting (*Helichrysum arenarium* L.): A Case of Stability Study", Plants, 2022

Publication

17

Sushma P. Ijardar, Vickramjeet Singh, Ramesh L. Gardas. "Revisiting the Physicochemical Properties and Applications of Deep Eutectic Solvents", *Molecules*, 2022

Publication

<1 %

18

archive-ouverte.unige.ch

Internet Source

<1 %

19

cwww.intechopen.com

Internet Source

<1 %

20

Raquel Cañadas, Blanca Sáenz de Miera, Paloma Méndez, Emilio J. González, María González-Miquel. "Enhanced Recovery of Natural Antioxidants from Grape Waste Using Natural Eutectic Solvents-Based Microwave-Assisted Extraction", *Molecules*, 2023

Publication

<1 %

21

cellulosicbiomasslab.engr.ucr.edu

Internet Source

<1 %

22

idoc.pub

Internet Source

<1 %

23

www.tandfonline.com

Internet Source

<1 %

24

link.springer.com

Internet Source

<1 %

25

Erika Dobroslavić, Maja Repajić, Verica Dragović-Uzelac, Ivona Elez Garofulić. "Isolation of Laurus nobilis Leaf Polyphenols: A Review on Current Techniques and Future Perspectives", *Foods*, 2022

Publication

<1 %

26

Harishbabu Balaraman, Rangabhashiyam Selvasembian, Vivek Rangarajan, Senthilkumar Rathnasamy. "Sustainable and Green Engineering Insights on Deep Eutectic Solvents toward the Extraction of Nutraceuticals", *ACS Sustainable Chemistry & Engineering*, 2021

Publication

<1 %

27

Khoddami, Ali, Meredith Wilkes, and Thomas Roberts. "Techniques for Analysis of Plant Phenolic Compounds", *Molecules*, 2013.

Publication

<1 %

28

Luis Carrillo-Hormaza, Luisa Duque, Sebastián López-Parra, Edison Osorio. "High-intensity ultrasound-assisted extraction of Garcinia madruno biflavonoids: mechanism, kinetics, and productivity", *Biochemical Engineering Journal*, 2020

Publication

<1 %

29 N. Sukor, R. Jusoh, S.A. Rahim, N. Kamarudin. <1 %
"Ultrasound assisted methods for enhanced
extraction of phenolic acids from Quercus
Infectoria galls", Materials Today:
Proceedings, 2018
Publication

30 Xianchao Shang, Jia-Neng Tan, Yongmei Du, <1 %
Xinmin Liu, Zhongfeng Zhang.
"Environmentally-Friendly Extraction of
Flavonoids from Cyclocarya paliurus (Batal.)
Ijinskaja Leaves with Deep Eutectic Solvents
and Evaluation of Their Antioxidant Activities",
Molecules, 2018
Publication

31 Yuntao Dai, Jaap van Spronsen, Geert-Jan <1 %
Witkamp, Robert Verpoorte, Young Hae Choi.
"Ionic Liquids and Deep Eutectic Solvents in
Natural Products Research: Mixtures of Solids
as Extraction Solvents", Journal of Natural
Products, 2013
Publication

32 pubs.rsc.org <1 %
Internet Source

33 res.mdpi.com <1 %
Internet Source

34 vdoc.pub <1 %
Internet Source

35

www.scielo.br

Internet Source

<1 %

36

"Application of Ionic Liquids in Biotechnology",
Springer Science and Business Media LLC,
2019

Publication

<1 %

37

Emanuelle de Faria, Rafael do Carmo, Ana
Filipa Cláudio, Carmen Freire, Mara Freire,
Armando Silvestre. "Deep Eutectic Solvents as
Efficient Media for the Extraction and
Recovery of Cynaropicrin from *Cynara
cardunculus* L. Leaves", *International Journal
of Molecular Sciences*, 2017

Publication

<1 %

38

Hamza Alasalvar, Zeliha Yildirim. "Ultrasound-
assisted extraction of antioxidant phenolic
compounds from *Lavandula angustifolia*
flowers using natural deep eutectic solvents:
An experimental design approach",
Sustainable Chemistry and Pharmacy, 2021

Publication

<1 %

39

Matteo Perra, Laura Fancello, Ines Castangia,
Mohamad Allaw et al. "Formulation and
Testing of Antioxidant and Protective Effect of
Hyalurosomes Loading Extract Rich in
Rosmarinic Acid Biotechnologically Produced
from *Lavandula angustifolia* Miller",
Molecules, 2022

<1 %

40

Saha, S.. "Some connectivity based cluster validity indices", Applied Soft Computing Journal, 201205

Publication

<1 %

41

Sujata S. Patil, Ajay Pathak, Virendra K. Rathod. "Optimization and Kinetic Study of Ultrasound assisted Deep Eutectic Solvent based extraction: A greener route for extraction of curcuminoids from Curcuma longa", Ultrasonics Sonochemistry, 2020

Publication

<1 %

42

Tao-Xiang Yang, Li-Qing Zhao, Juan Wang, Guo-Li Song, Hai-Min Liu, Hui Cheng, Zhen Yang. "Improving Whole-Cell Biocatalysis by Addition of Deep Eutectic Solvents and Natural Deep Eutectic Solvents", ACS Sustainable Chemistry & Engineering, 2017

Publication

<1 %

43

archive.org

Internet Source

<1 %

44

dokumen.pub

Internet Source

<1 %

45

encyclopedia.pub

Internet Source

<1 %

46

uest.ntua.gr

Internet Source

<1 %

47

www.kgshome.org

Internet Source

<1 %

48

Kavita J. Lanjekar, Virendra K. Rathod. " Application of Ultrasound and Natural Deep Eutectic Solvent for the Extraction of Glycyrrhizic Acid from : Optimization and Kinetic Evaluation ", Industrial & Engineering Chemistry Research, 2021

Publication

<1 %

49

Sujata S. Patil, Siddhant Bhasarkar, Virendra K. Rathod. " Extraction of curcuminoids from : comparative study between batch extraction and novel three phase partitioning ", Preparative Biochemistry and Biotechnology, 2019

Publication

<1 %

50

Nurmayulis, F R Eris, D Hastuti, R T Utami, Y R Denny, T Firmansyah. " Effects of neem plant extract (A. Juss) and bio-surfactant diethanolamide olien from palm oil to the mortality of cacao moth pest () ", IOP Conference Series: Earth and Environmental Science, 2019

Publication

<1 %

51

Oscar Zannou, Hojjat Pashazadeh, Mohamed Ghellam, Salam A. Ibrahim, Ilkay Koca. "Extraction of Anthocyanins from Borage (Echium amoenum) Flowers Using Choline

<1 %

Chloride and a Glycerol-Based, Deep Eutectic Solvent: Optimization, Antioxidant Activity, and In Vitro Bioavailability", *Molecules*, 2021

Publication

52

Wei Wang, Xiaoli Zhang, Qinglei Liu, Yucheng Lin, Zhiguo Zhang, Shanshan Li. "Study on Extraction and Antioxidant Activity of Flavonoids from *Hemerocallis fulva* (Daylily) Leaves", *Molecules*, 2022

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On