

biodiversitas 2021

by Achmad Noerkhaerin Putra

Submission date: 02-Oct-2021 08:15AM (UTC+0700)

Submission ID: 1663003790

File name: 3._Article.pdf (632.11K)

Word count: 8610

Character count: 45107

Dietary *Bacillus* NP5 supplement impacts on growth, nutrient digestibility, immune response, and resistance to *Aeromonas hydrophila* infection of African catfish, *Clarias gariepinus*

ACHMAD NOERKHAERIN PUTRA^{1,2,*}, MUSTAHAL^{1,2}, MAS BAYU SYAMSUNARNO¹, DODI HERMAWAN¹, DEVIA GUSNUR FATIMAH¹, PRAMODITA BALITA PUTRI¹, SEVIA¹, RINA ISNAINI¹, MUHAMAD HERJAYANTO¹

¹Department of Fisheries, Faculty of Agriculture, Universitas Sultan Ageng Tirtayasa. Jl. Raya Jakarta Km. 04, Pakupatan, Serang 42121, Banten, Indonesia. Tel.: +62-254-280330, Fax.: +62-254-281254, *email: Putra.achmadnp@untirta.ac.id

²Indonesia-Center of Excellence for Local Food Innovation, Universitas Sultan Ageng Tirtayasa. Jl. Raya Jakarta Km. 04, Pakupatan, Serang 42121, Banten, Indonesia

Manuscript received: 9 October 2020. Revision accepted: 21 December 2020.

Abstract. Putra AN, Mustahal, Syamsunarno MB, Hermawan D, Fatimah DG, Putri PB, Sevia, Isnaini R, Herjayanto M. 2020. Dietary *Bacillus* NP5 supplement impacts on growth, nutrient digestibility, immune response, and resistance to *Aeromonas hydrophila* infection of African Catfish, *Clarias gariepinus*. *Biodiversitas* 22: 253-261. This study aims to investigate the effects of *Bacillus* NP5 supplementation as a probiotic on growth, immune response, and resistance of African catfish to *Aeromonas hydrophila* infection. Catfish with an initial weight of 6.8 ± 0.1 g were fed with different doses of *Bacillus* NP5 (0%, 1.1%, 1.2%, 1.3%, and 1.4%) with three replications and reared for 60 days. The first 45 days were used to observed growth performance, and the last 15 days were used to perform challenge tests against *A. hydrophila* infection. The results showed that the treatment of 1.2% *Bacillus* NP5 results in the highest specific growth rate (2.55 ± 0.28 day⁻¹) and increased protein and lipid digestibility significantly ($P < 0.05$). Treatment of 1.1-1.3% *Bacillus* NP5 increase amylase and lipase activity that significantly higher than 0% *Bacillus* NP5 treatment. Supplementation of *Bacillus* NP5 significantly increased the leukocyte, phagocytic index, and survival rate in African catfish after *A. hydrophila* infection. Therefore, the supplementation of 1.2% *Bacillus* NP5 in the feed increased the growth, immune response, and African catfish resistance to the infection.

Keywords: African catfish, *Bacillus* NP5, growth, probiotic, response immune

INTRODUCTION

Indonesia is the second-largest producer of aquaculture products in the world after China, with a total of 9.3 million tons in 2018 (FAO 2020). African catfish (*Clarias gariepinus*) is one of the common commodities found in its freshwater fishery. According to the Ministry of Marine Affairs and Fisheries (2019), catfish production in Indonesia was the 3rd largest after shrimp and tilapia, with a total production of 19.6 thousand tons in 2017. Catfish farming has been intensively and sustainably carried out to fulfill the increasing demand for fish. Zhang et al. (2018) stated that various negative impacts, such as the emergence of disease attacks, decrease in environmental quality, and retardation of fish growth, are associated with the intensification of aquaculture. *Aeromonas hydrophila* is an opportunistic gram-negative bacterium. It causes Motile *Aeromonas* Septicemia (MAS) in fish (Korni and Khalil 2017; Shoemaker et al. 2018), which leads to an increase in mortality rate (Maji et al. 2017; Jung-Schroers et al. 2018; Li et al. 2019a). Over the past few decades, fish farmers have adopted the use of antibiotics, a conventional method to control disease attacks (Lee et al. 2019). However, antibiotics and other chemical compounds lead to the improvement of resistance in pathogenic bacteria (Hoseinifar et al. 2018) and increase chemical residues in

water and fish (Lee et al. 2016; Hassani et al. 2019). Therefore, an environmentally friendly approach is needed to overcome the problems in aquaculture intensification to maintain sustainability.

Probiotics are environmentally friendly applications to improve fish health and growth (Carnevali et al. 2017). According to Ramesh et al. (2015), it is a substitute for antibiotics and can suppress pathogens' growth without being detrimental to the host and its environment. Furthermore, Li et al. (2019b) reported that it has been widely applied in aquaculture and used as a substitute for fish and shrimp antibiotics. Previous studies showed that probiotics supplementation in aquaculture results in increasing growth and feed digestibility. Probiotics increase digestive enzyme activity (Enferadi et al. 2018; Jallahi et al. 2018; Arani et al. 2019), increase immunity against pathogens (Amir et al. 2019; Zhu et al. 2019) and improve the water quality of culture media (Elsabagh et al. 2018; Kewcharoen and Srisapoome 2019).

Bacillus NP5 is a Gram-positive bacteria, oxidation fermentation, and catalase-positive (Putra and Widanarni 2015). Previous studies showed that supplementation of *Bacillus* NP5 in fish increase growth (Putra et al. 2015; Utami et al. 2015), increase immunity against fish pathogens (Widanarni et al. 2014; Febrianti et al. 2016; Tamamdusturi et al. 2016) and improve water quality

(Putra et al. 2020a). The addition of 1% and 1.52% of *Bacillus* NP5 results in increasing African catfish' growth performance (Putra et al. 2020b). In this study, a probiotic dose range of 1-1.5% was used to obtain the correct dose of *B. NP5* in the diet. Besides, there were still a few data related to the effect of *Bacillus* NP5 addition to catfish's immune system. Therefore, this study was carried out to investigate *Bacillus* NP5 probiotics' supplementation on growth performance, immune response, and African catfish resistance to *A. hydrophila* infection.

MATERIALS AND METHODS

Probiotic and feed preparation

Bacillus NP5 is a probiotic bacteria originating from the intestine of *Oreochromis niloticus* (Putra and Widanarni 2015). Bacteria were cultured according to the method by Putra and Romdhonah (2019). *Bacillus* NP5 was cultured aseptically in 250 mL of tryptic soy broth (TSB) medium and incubated for 18 hours at room temperature. After incubation, the bacterial culture was centrifuged (HITACHI himac CT6E/CT6EL) at the speed of 1000 x g for 10 minutes at 4°C, and the bacterial supernatant was washed and homogenized by adding 100 mL of phosphate buffer saline (PBS, pH 7.2). Furthermore, serial dilutions with PBS were carried out to obtain a 10⁶ CFU/mL bacterial density. Bacterial supernatant was stored at 4°C before being mixed in the feed.

Commercial catfish feed (781, INDONESIA) was used as a basal diet grounded into flour, sieved with 320 µm mesh, and thoroughly mixed with chromium oxide of 0.5% as the digestibility indicator and 3% tapioca as a binder. Then, 30% of warm water was added and printed in a 1-2 mm feed diameter in a pellet machine. Subsequently, the feed was oven heated at 50°C for 24 hours, followed by homogenization of 2% egg yolk (Putra et al. 2015) and sprayed with *Bacillus* NP5 supernatant according to the treatment dose. The proximate composition of the experimental diet is presented in Table 1.

Table 1. Proximate composition (g/100 g feed) of experimental diet

Proximate	Value
Crude protein (%)	33.95
Crude lipid (%)	7.73
Crude fiber (%)	4.16
Ash (%)	9.9
Moisture (%)	6.49
Organic matter (%)*	83.61
Nitrogen-free extracts**	37.77
Dry matter (%)	93.51
Energy (Kcal g feed ⁻¹ ***)	275.863
Size feed (mm)	1-2

Note: *Dry weight - ash, **100 - protein-lipid - fiber - ash - moisture, **Digestible energy according to NRC (1993), calculated as protein 118.8 Kcal g feed⁻¹; lipid 62.6 Kcal g feed⁻¹, and carbohydrate: 94.4 Kcal g feed⁻¹

Fish rearing and experimental procedure

A total of 375 African catfish reared from the Baros Seed Center, Serang, Indonesia, were used in this study. Fish rearing was carried out at the Laboratory of Aquaculture, University of Sultan Ageng Tirtayasa, Indonesia. The initial weight of the catfish was 6.8 ± 0.1 g were randomly scattered into the round tank with a diameter of 72 cm diameter and fish density of 25 fish /tank with a water volume of 40 L. The aerator was placed at the center of the tank at 15 cm from the bottom. The acclimatization process was carried out for seven days, and the fish was unfed for 24 hours before treatment. Feeding was carried out three times daily, i.e., at 08.00, 12.00, and 17.00 Western Indonesian time zone through a satiation method. Approximately 20% of the total water volume was changed every three days and 50% every two weeks. The rearing process was carried out for 60 days, of which the first 45 days were used for growth tests and the remaining 15 days for a challenge test with the injection of the pathogen *A. hydrophila*.

A completely randomized design with five different levels of *Bacillus* NP5 probiotic (0% or negative control, 1.1%, 1.2%, 1.3%, and 1.4%) with three replications was used in this study. The in-situ water quality monitoring such as temperature (RESUN, Indonesia) was carried out daily, while pH (LUTRON 208) and dissolved oxygen (LUTRON DO550) were observed weekly. Ammonia measurements were carried out based on the standard Phenate method (APHA 1998) at the beginning and the end of the rearing period. In the present study, water temperature was in the range of 27.1-28.8°C, while the pH value during rearing before and after water change was 5.17-7.23. The dissolved oxygen was 5.17-7.23 mg L⁻¹, while the ammonia value was below 2 mg L⁻¹. The value of water quality showed that it was in the normal range for catfish farming, according to Bhatnagar and Devi (2013).

Growth parameters

The number and weight of African catfish were measured at the beginning and the 45 days to determine the growth performance, which was calculated based on the equation proposed by Huisman (1987) as follow: (i) Feed intake (g) = the initial weight of feed - final weight of feed. (ii) Specific growth rate (% day⁻¹) = 100 × ((ln final weight of catfish - ln initial weight of catfish / 50 days)). (iii) Feed conversion ratio = Total feed consumption / final weight of catfish - the initial weight of catfish. (iv) Survival rate (%) = 100 × (final number of catfish / initial number of catfish)

Collection of feces was carried out 2 hours after feeding by siphon from the 7th day until the treatment was complete. Feces was stored at -4°C. Nutrient digestibility was calculated based on the equation proposed by Takeuchi (1988) as follows: (i) Dry matter digestibility (%) = 100 × (1 - (% chromium content in the diet / % chromium content in the feces)). (ii) Protein/lipid digestibility (%) = 100 × (1 - ((% chromium content in the diet / % chromium content in the feces) × (% nutrient content in the feces / % nutrient content in the feed)))

Digestive enzyme activity was determined on day 45. A total of 5 fish from each treatment were randomly selected, and their total intestines (anterior and posterior) were surgically removed and weighed. The intestinal samples were crushed to obtain the substrate. The Bergmeyer et al. (1983) method was used to determine the protease activity with casein as a substrate in 0.05 M phosphate buffer, pH 7, and 2 mmol/L of tyrosine. Amylase activity was determined using 1% starch solution as a substrate in 20 mM sodium phosphate buffer, pH 6.9, and 20 mM of sodium phosphate according to the method by Worthington (1993). Lipid enzyme activity was calculated based on the procedure described by Borlongan (1990), using olive oil as a substrate and Tris-HCL as a buffer.

Chemical analysis

Proximate analysis of feed and fish was carried out using the standard method of AOAC (2005). A total of 5 catfish were randomly selected for analysis with the micro-Kjeldahl method to determine crude protein content. Moisture was estimated with an oven at 105°C for 6 hours, while the sample was analyzed through extraction with a Soxhlet to determine crude lipid content. The percentage of ash was determined by combustion at 550 °C for 24 hours, and crude fiber was estimated by an Automatic analyzer.

Experimental infection

Aeromonas hydrophila was obtained from the Fish Health Laboratory of IPB University for challenge tests carried out on the 45th day. Before the challenge test, Koch's Postulates were performed to increase the virulence of *A. hydrophila*. Catfish were reared according to the treatment, and the fish density was uniform (15 individuals/tank). Meanwhile, *A. hydrophila* were cultured on 200 ml of TSB media for 24 hours and incubated at 29 °C. After incubation, the culture was centrifuged at 1000 x g for 10 minutes and diluted using PBS (pH: 7.2) to obtain 10⁷ CFU/mL density. Furthermore, it was injected intramuscularly into catfish as much as 0.1 mL per fish (Putra and Widanarni 2015).

Hematological and immune response

Evaluations of the hematological and immune response were carried out on days 0, 45, and 60. Five fish from each treatment were randomly selected to determine the number of erythrocytes and leukocytes according to the method by Blaxhall and Daisley (1973). Blood was mixed with Hayem's and Turk's solutions until it was homogeneous. Erythrocytes were counted using a Neubauer-type hemocytometer under a microscope with a magnification of 400x. Hematocrit level was calculated using a microhematocrit tube following the method by Anderson and Siwicki (1995) Determination of hemoglobin levels in the blood refers to Sahli's method (Wedemeyer and Yasutake 1977). Phagocytosis index was measured by mixing 20 µL of the blood sample with 20 µL of *A.*

hydrophila suspension as an antigen with a density of 10⁷ cells mL⁻¹. The phagocyte cells are counted based on the percentage during the phagocytosis process (Ispir and Dorucu 2005).

Statistical analysis

Data of growth and immune response were analyzed by Analysis of Variance (ANOVA) in Microsoft Office Professional Plus 2010. Duncan Multiple Range Test (DMRT) analyzed significant differences between treatments with a 95% confidence interval.

RESULTS AND DISCUSSION

Nutrient digestibility and digestive enzyme activity of *A. hydrophila* in catfish

Table 2 showed that the dry matter digestibility in treatment 1.1-1.2% *Bacillus NP5* was significantly higher ($P < 0.05$) compared to 0% *Bacillus NP5*. However, the dry matter digestibility in the treatment of 1.3-1.4% *Bacillus NP5* was not significantly different compared to the control (0%) ($P > 0.05$). Treatment of 0% and 1.3% *Bacillus NP5* results in the lowest protein digestibility, while the highest was in 1.2% *Bacillus NP5* ($P < 0.05$) treatment. The lowest lipid digestibility was obtained from 0% and 1.4% *Bacillus NP5* treatment and the highest one was in 1.1-1.3% *Bacillus NP5* ($P < 0.05$). Furthermore, the highest protease enzyme activity was obtained from 1.2% *Bacillus NP5*, significantly different from other treatments ($P < 0.05$). The highest amylase and lipase enzyme activity were obtained from 1.1-1.3% *Bacillus NP5* treatment.

Growth performance

The growth performance of catfish was presented in Table 3. The highest specific growth rate was in 1.2% *Bacillus NP5* treatment, which significantly different from other treatments ($P < 0.05$), while treatments of 0%, 1.1%, 1.3%, and 1.4% *Bacillus NP5* were not different. The final weights of 1.1-1.3% *Bacillus NP5* treatments were significantly higher than 0% *Bacillus NP5*. However, feed intake and survival rates in all treatments were not significantly different ($P > 0.05$). Furthermore, the highest feed conversion was obtained from 0% *Bacillus NP5* treatment, which was significantly different from the treatment of 1.1-1.4% *Bacillus NP5* ($P < 0.05$).

Whole-body composition

The whole-body composition of catfish was presented in Table 4. The highest crude protein was obtained from 1.2% *Bacillus NP5* treatment ($P < 0.05$). Crude lipids in treatments of 1.2-1.4% *Bacillus NP5* were significantly higher than 0-1.1% *Bacillus NP5* treatment. However, the application of *Bacillus NP5* does not have any effect on moisture and ash content ($P > 0.05$).

Table 2. Nutrient digestibility and digestive enzyme activity of African catfish with five experimental diets containing the different doses of *Bacillus* NP5

Dietary of probiotic	Digestibility and digestive enzyme parameters*					
	Dry matter digestibility (%)	Protein digestibility (%)	Lipid digestibility (%)	Protease activity (U. mg protein ⁻¹)	Amylase activity (U. mg protein ⁻¹)	Lipase activity (U. mg protein ⁻¹)
0% <i>B. NP5</i>	66.86±0.97 ^a	76.06±1.31 ^a	56.85±4.39 ^a	0.007±0.00 ^a	1.40 ±0.04 ^a	0.282±0.00 ^a
1.1% <i>B. NP5</i>	69.09±1.70 ^b	78.10±1.11 ^c	67.75±0.94 ^{bc}	0.014±0.00 ^b	2.07±0.034 ^b	0.325±0.016 ^{bc}
1.2% <i>B. NP5</i>	73.01±1.27 ^c	85.52±1.14 ^d	70.30±2.34 ^c	0.057±0.00 ^d	2.38 ±0.05 ^b	0.358±0.00 ^c
1.3% <i>B. NP5</i>	67.75±0.94 ^{ab}	76.16±0.47 ^{ab}	70.86±0.45 ^c	0.037±0.00 ^c	2.35 ±0.08 ^b	0.354±0.03 ^c
1.4% <i>B. NP5</i>	77.78±0.33 ^a	77.39±0.72 ^{bc}	62.32±6.46 ^{ab}	0.010±0.00 ^a	1.28±0.26 ^a	0.308±0.01 ^{ab}

Note: ^a mean ± SD values in the same column with similar superscript letters are not significantly different ($P > 0.05$)

Table 3. Growth performance of African catfish with five experimental diets containing the different doses of *Bacillus* NP5

Dietary of probiotic	Growth parameters*					
	Initial weight (g fish ⁻¹)	Final weight (g fish ⁻¹)	Feed intake (g)	Specific growth rate (% day ⁻¹)	Feed conversion ratio	Survival rate (%)
0% <i>B. NP5</i>	6.77±0.02	19.53 ± 1.18 ^a	321.37±57.62	2.12±0.12 ^a	1.82±0.07 ^c	96.00± 0.00
1.1% <i>B. NP5</i>	6.74±0.02	23.78 ± 2.89 ^{bc}	330.37±26.75	2.16±0.14 ^a	1.29±0.02 ^b	94.67±2.30
1.2% <i>B. NP5</i>	6.75±0.01	26.26 ± 1.14 ^c	328.33±9.28	2.55±0.28 ^b	1.15±0.04 ^a	96.00 ± 0.00
1.3% <i>B. NP5</i>	6.84±0.26	23.91 ± 1.34 ^{bc}	329.62±8.85	2.35±0.10 ^{ab}	1.17±0.02 ^a	96.00±4.00
1.4% <i>B. NP5</i>	6.99±0.01	21.98 ± 1.01 ^{ab}	324.93±28.55	2.34±0.06 ^{ab}	1.28±0.02 ^b	97.33 ±2.30

* mean ± SD values in the same column with similar superscript letters are not significantly different ($P > 0.05$)

Table 4. Whole-body composition of African catfish with five experimental diets containing different doses of *Bacillus* NP5

Dietary of probiotic	Body composition*			
	Crude protein (%)	Crude lipid (%)	Moisture (%)	Ash (%)
0% <i>B. NP5</i>	10.35±0.05 ^a	5.01±0.02 ^a	71.07±0.46	2.82±0.05
1.1% <i>B. NP5</i>	10.07±0.03 ^a	5.11±0.09 ^a	71.15±0.02	2.80±0.12
1.2% <i>B. NP5</i>	13.16±0.28 ^c	5.52±0.10 ^b	72.66±0.35	2.86±0.03
1.3% <i>B. NP5</i>	11.52±0.85 ^b	5.54±0.33 ^b	73.96±0.40	2.77±0.06
1.4% <i>B. NP5</i>	12.05±1.12 ^{bc}	5.12±0.10 ^b	73.36±0.98	2.87±0.07

* mean ± SD values in the same column with similar superscript letters are not significantly different ($P > 0.05$)

Hematological and immune response

Table 5 showed no significant difference in the values of blood profile and phagocyte index of African catfish on day 0. The same results were also obtained on day 45, except for the hematocrit value in the probiotic treatment, which was higher than the control ($P < 0.05$). At the end of rearing (day 60), the highest hemoglobin value was obtained from 1.2% *Bacillus* NP5, which was higher than other treatments ($P < 0.05$). Erythrocyte value in the treatment of 1.1, 1.2, and 1.4% *Bacillus* NP5 was not significantly different, but it significantly higher than the 0% *Bacillus* NP5 ($P < 0.05$) treatment. Probiotic treatment had a higher value of hematocrit than 0% *Bacillus* NP5 ($P < 0.05$). Supplementation of 1.1-1.2% *Bacillus* NP5 results in higher leukocytes than other treatments ($P < 0.05$). The highest phagocytes index was obtained from 1.2% *Bacillus* NP5 ($P < 0.05$). The lowest survival rate was at 0% *Bacillus* NP5 treatment in weeks 1 and 2 (Figure 1). The highest survival rate after *A. hydrophila* infection was obtained from 1.2% *Bacillus* NP5 treatment, namely 75.46% and 66.67% in weeks 1 and 2, respectively.

Discussion

Bacteria of the genus *Bacillus* have been used as probiotics in aquaculture with numerous beneficial effects on their hosts, such as increasing growth performance, enhancing feed digestibility, the production of exogenous enzymes, enhancing immunity against pathogens, and stimulating microbiota populations to inhibit pathogenic colonization (Kuebutornye et al. 2019; Ringø 2020). The high activity of digestive enzymes indicates an increase in the development of the intestine of fish and its ability to digest feed (Darafsh et al. 2019). According to Iqbal and Khosio (2016), protease, lipase, and amylase are the main enzymes responsible for feed digestion in the digestive tract of fish. This study showed that *Bacillus* NP5 supplementation increased protease activity, amylase, and lipase in the intestine of catfish compared to controls, especially at doses of 1.1-1.3%. It increases is due to the production of exoenzyme by probiotics. Sankar et al. (2016) reported that the high activity of digestive enzymes in feed treated with probiotics is due to the exoenzyme production by probiotics and its stimulation to the digestive tract of the host. The results obtained in this research are in

line with the previous studies by Putra et al. (2015 and 2021) that the administration of *Bacillus NP5* increases the activity of protease, lipase, and amylase enzymes in the digestive tract of the host. Similar results were found on *L. vannamei* (Zokaeifar et al. 2012; Zokaeifar et al. 2014), Javanese carp (Allameh et al. 2015), and tilapia (Midhun et al. 2019).

The results showed that supplementation of 1.1-1.2% *Bacillus NP5* increased nutrient stability compared to 0% *Bacillus NP5*. A study by Lin et al. (2004) showed that supplementation of *Bacillus* spp. increased the digestibility of dry matter and protein in white shrimp. Mohapatra et al. (2012) reported the same results, which showed that dietary

probiotics in feed significantly ($P < 0.05$) enhance dry matter, protein, and lipid digestibility compared to controls on the rearing of juvenile rohu, *Labeo rohita*. In this study, the supplementation of 1.2% *Bacillus NP5* significantly increased the digestibility of dry matter and protein ($P < 0.05$), which were considered to be related to the protease enzyme activity in 1.2% *Bacillus NP5* treatment. Meanwhile, increasing protease enzyme activity in probiotic treatment has been reported in other fish species, such as *Litopenaeus vannamei*, with the application of *Lactobacillus pentosus* AS13 (Zheng and Wang 2016) and rainbow trout with *L. plantarum* supplementation (Enferadi et al. 2018).

Table 5. The hematological and immune response of African catfish with five experimental diets containing the different doses of *Bacillus NP5*

Parameter/day	Dietary of probiotic*				
	0% <i>Bacillus NP5</i>	1.1% <i>Bacillus NP5</i>	1.2% <i>Bacillus NP5</i>	1.3% <i>Bacillus NP5</i>	1.4% <i>Bacillus NP5</i>
0 day					
Hemoglobin (g%)	6.20±0.00	6.23±0.12	6.24±0.70 ^{ab}	6.21±0.9	6.21±0.69
Erythrocytes (x 10 ⁶ cell/mm ³)	2.89±0.09	2.85±0.04	2.95±0.05	2.81±0.08	2.81±0.08
Hematocrit (%)	17.88±0.3	17.60±0.17	17.96±0.30	20.17±2.49	20.17±2.49
Leukocytes (x 10 ⁴ cell/mm ³)	6.32±0.02	6.35±0.06	6.34±0.03	6.16±0.36	6.16±0.36
Phagocyte index (g%)	40.33±4.51	40.33±4.73	40.00±4.36	40.07±1.53	40.67±1.53
45 days**					
Haemoglobin (g%)	7.03±0.42	6.94±0.53	7.00±0.20	7.09±0.62	6.95±0.64
Erythrocytes (x 10 ⁶ cell/mm ³)	2.16± 0.14	2.60±0.55	2.70±0.02	2.38±0.11	2.66±0.33
Hematocrit (%)	7.82±0.64 ^a	11.27±0.37 ^b	11.24±0.69 ^b	11.36±1.29 ^b	11.36±1.16 ^b
Leukocytes (x 10 ⁴ cell/mm ³)	8.74±0.02	8.63±0.02	8.95±0.03	8.49±0.28	8.86±0.22
Phagocyte index (g%)	41.67±4.04	40.67±4.51	40.67±7.37	40.00±3.46	41.33±4.51
60 days***					
Hemoglobin (g%)	4.40±0.40 ^a	5.60±0.72 ^{ab}	6.00±0.20 ^b	5.26±1.16 ^{ab}	5.38±1.22 ^{ab}
Erythrocytes (x 10 ⁶ cell/mm ³)	1.25±0.04 ^a	1.81±0.03 ^b	1.45±0.05 ^b	1.35±0.09 ^{ab}	1.45±0.05 ^b
Hematocrit (%)	7.80±0.40 ^a	14.20±0.25 ^b	12.56±0.39 ^b	12.32±0.30 ^b	15.40±5.90 ^b
Leukocytes (x 10 ⁴ cell/mm ³)	8.49±0.12 ^a	9.54±0.02 ^b	9.45±0.03 ^b	9.53±0.50 ^b	9.40±0.39 ^b
Phagocyte index (g%)	51.33±2.52 ^a	46.67±1.53 ^a	66.00±3.00 ^b	62.67±5.03 ^b	58.67±4.73 ^{ab}

Note: ^amean ± SD values in the same row with similar superscript letters are not significantly different ($P > 0.05$). **before *Aeromonas hydrophila* infection, ***after *A. hydrophila* infection.




Figure 1. The survival rate of African catfish after *Aeromonas hydrophila* infection

The application of probiotics in feed has resulted in the increased growth of fish and shrimp (Liu et al. 2017; Arani et al. 2019; Xie et al. 2019). Jang et al. (2019) stated that probiotics increase the growth of the host by improving the digestibility of the feed. The highest final weight, specific growth rate, and feed conversion ratio were obtained from 1.2% *Bacillus* NP5 treatment ($P < 0.05$) was considered to be associated with high nutrient digestibility due to increased enzyme activity. According to Wu et al. (2012), probiotic supplementation improves the digestion process of feed and the beneficial microbial population and digestive enzyme activity in the digestive tract of the host. These results were consistent with previous studies, which showed that probiotics have a positive effect on enzyme activity, feed digestibility, and growth performance in hybrid groupers (He et al. 2016), tilapia (Poolsawat et al. 2020), and *L. vannamei* (Tsai et al. 2010).

The results showed that the feed intake was the same for all treatments, indicating that *Bacillus* NP5 supplementation in the feed did not affect its palatability (Tantikitti et al. 2014). Similar results were observed in tilapia (Elsabagh et al. 2018), juvenile olive flounder (Lee et al. 2019), and rainbow trout (Nargesi et al. 2019). Supplementation of 1.2-1.4% *Bacillus* NP5 increased crude protein and lipids in the body composition of African catfish. It was consistent with the results obtained in several other species, such as goldfish, by administering *L. delbrueckii* (Zhang et al. 2018) and *L. vannamei* through the use of *B. subtilis* and *B. licheniformis* (Madani et al. 2018). Increasing protein and lipids in the body composition can be caused by the increased digestibility and food absorption of feed. As a result, there were more protein and fat deposits in the body (Abdel-Tawwab et al. 2008; Reda and Selim 2015). The results showed no difference in moisture and ash content in all treatments ($P > 0.05$). Similar results were discovered in *L. vannamei* (Adel et al. 2017), parrotfish (Liu et al. 2017), and snakehead (Dai et al. 2018).

Several studies have reported that probiotic supplementation reduces the mortality rate of fish and improves its immune system against pathogens (Adel et al. 2017; He et al. 2017; Hassani et al. 2019). Dawood et al. (2016) stated that probiotic supplementations in feed positively affect fish blood profile. This study showed that the erythrocyte and hemoglobin number in African catfish decreased after the challenge test compared to the pre-challenge test. Olivier et al. (1981) stated that the decrease in erythrocyte and hemoglobin is related to the ability of pathogens to damage red blood cells. A study by Kumar et al. (2015) showed that the number of erythrocytes and hemoglobin in Catla fingerling after experiencing a challenge test with *A. hydrophila* bacteria was decreased. The hematocrit number was the ratio of erythrocyte to total blood volume influenced by the number of cells (Soltani et al. 2017). In this study, the hematocrit value obtained by probiotic treatment was higher before and after pathogenic bacterial infection compared to those without probiotics. This result indicates that probiotic has improved the

immune system of African catfish. Hematocrit value is related to health status of fish (Nandi et al. 2017) and administration probiotics stimulated the innate immunity of fish (Adorian et al. 2018). The same results demonstrated by Ahmadifar et al. (2020) who reported the addition of probiotic *Pediococcus pentosaceus* 1×10^9 CFU g^{-1} increased the general hematocrit value compared to the controls in carp. Zhu et al. (2019) also reported that the hematocrit value in tilapia given probiotic *B. subtilis* LT3-1 was increased compared to the controls.

In this study, *Bacillus* NP5 supplementation in feed significantly increased leukocytes after infection with *A. hydrophila*. These results were in line with the study carried out by Reda and Selim (2015) that supplementation of *B. amyloliquefaciens* increased the leukocyte of tilapia. A study by Pourgholam et al. (2017) showed that the addition of *L. plantarum* as a probiotic increased the leukocyte number in juvenile Siberian sturgeon. The leukocyte was one of the fish's non-specific defense systems (Uribe et al. 2011). The increase in fish leukocytes was related to the body's defense system response to disease infections (Tanbiyaskur et al. 2015). Phagocytes have been widely used to determine the defense against pathogenic infections (Giri et al. 2012). Tamamdusturi et al. (2016) stated that phagocytosis is the initial cellular immune system after a pathogenic infection was provided by monocytes and granulocytes. According to Djauhari et al. (2016), probiotic supplementation increases the cellular immune system of fish and increasing its resistance to pathogenic infections. The phagocyte index of African catfish increased significantly in treatment of *Bacillus* NP5 after the challenge test. This result was similar to the result of a study by Doan et al. (2015) on *Pangasius* catfish that was challenged with *A. hydrophila* and Zhao et al. (2019) on giant freshwater prawns given *B. pumilus* probiotic.

The increase in blood profile values, especially the leukocytes and phagocyte index in probiotic treatment, indicated an increase in the immune system of African catfish. It was confirmed by increasing survival rate after the challenge test with *A. hydrophila* at the *Bacillus* NP5 treatments. The results showed that *Bacillus* NP5 supplementation increased African catfish resistance to *A. hydrophila* infection and increased their survival rate compared to treatment without probiotics. It implies that *Bacillus* NP5 supplementation increases the cellular and humoral defenses of African catfish. Similar results were also reported by Zokaeifar et al. (2014) with the addition of *B. subtilis* in the feed, which was used to increase the survival rate of *L. vannamei* infected with *V. harveyi*. This is also in accordance with the result in freshwater prawn through the addition of *B. licheniformis* in the feed (Kumar et al. 2013), *Cyprinus carpio* with administration *L. delbrueckii* (Zhang et al. 2017), and white shrimp with *Pediococcus pentosaceus* supplementation in the diet (Adel et al. 2017).

In conclusion, dietary *Bacillus* NP5 supplement positively increased the growth, immune response, and resistance of African catfish against *A. hydrophila* infection

with the optimal dose of *Bacillus NP5* supplementation in the feed of African catfish was 1.2%.

ACKNOWLEDGEMENTS

The authors express gratitude to LPPM-UNTIRTA for funding this study through the research internal grant scheme of Penelitian Terapan Internal 2020 (Nomor: B/135/UN.43.9/PT.01.03/2020).

REFERENCES

- Abdel-Tawwab M, Abdel-Rahman AM, Ismael NEM. 2008. Evaluation of commercial live bakers' yeast, *Saccharomyces cerevisiae* as a growth and immunity promoter for fry Nile tilapia, *Oreochromis niloticus* (L.) challenged *in situ* with *Aeromonas hydrophila*. *Aquaculture* 280: 185-189. DOI: 10.1016/j.aquaculture.2008.03.055.
- Adorian TJ, Jamali H, Farsani HG, Darvishi P, Hasanpour S, Bagheri T, Roozbehfar R. 2018. Effect of probiotic bacteria *Bacillus* on growth performance, digestive enzyme activity, and hematological parameters of Asian sea bass, *Lates calcarifer* (Bloch). *Probiotics Antimicrob Proteins* 11: 248-255. DOI: 10.1007/s12602-018-9393-z.
- Adel M, Yeganeh S, Dawood MAO, Safari R, Radhakrishnan S. 2017. Effects of *Pediococcus pentosaceus* supplementation on growth performance, intestinal microflora and disease resistance of white shrimp, *Litopenaeus vannamei*. *Aquac Nutr* 23 (6): 1401-1409. DOI: 10.1111/anu.12515.
- Ahmadifar E, Sadegh TH, Dawood MAO, Dadar M, Sheikhzadeh N. 2020. The effects of dietary *Pediococcus pentosaceus* on growth performance, hemato-immunological parameters and digestive enzyme activities of common carp (*Cyprinus carpio*). *Aquaculture* 516: 734656. DOI: 10.1016/j.aquaculture.2019.734656.
- Allameh SK, Ringø E, Yusoff FM, Daud HM, Ideris A. 2015. Dietary supplement of *Enterococcus faecalis* on digestive enzyme activities, short-chain fatty acid production, immune system response and disease resistance of Javanese carp (*Puntius gonionotus*, Bleeker 1850). *Aquac Nutr* 23 (2): 331-338. DOI: 10.1111/anu.12397.
- Amir I, Zuberi A, Kamran M, Imran M, Murtaza MH. 2019. Evaluation of commercial application of dietary encapsulated probiotic (*Geotrichum candidum* QAUGC01): Effect on growth and immunological indices of rohu (*Labeo rohita*, Hamilton 1822) in semi-intensive culture system. *Fish Shellfish Immunol* 95: 464-472. DOI: 10.1016/j.fsi.2019.11.011.
- Anderson DP, Siwicki AK. 1995. Basic hematology and serology for fish health program. *Proceeding of the Second Symposium on Disease in Asian Aquaculture*, October 25-29, 1995.
- Association of Official Analytical Chemists (AOAC). 2005. *Official Methods of Analysis*, 16th ed. AOAC, Arlington, VA, USA.
- Arani MM, Salati AP, Safari O, Keyvanshokoo S. 2019. Dietary supplementation effects of *Pediococcus acidilactici* as probiotic on growth performance, digestive enzyme activities, and immunity response in zebrafish (*Danio rerio*). *Aquac Nutr* 25 (4): 854-861. DOI: 10.1111/anu.12904.
- APHA. 1998. *Standard Methods for the Examination of the Water and Wastewater* (22nd ed.). American Public Health Association, Washington DC, USA.
- Bergmeyer HU, Grossl M, Walter HE. 1983. Reagents for enzymatic analysis. In: Bergmeyer HU (eds.). *Methods in Enzymatic Analysis*. Verlag Chemie, Weinheim.
- Bhatnagar A, Devi P. 2013. Water quality guidelines for the management of pond fish culture. *Intl J Environ Sci* 3 (6): 1980-2009. DOI: 10.6088/ijes.2013030600019.
- Blaxhall PC, Daisley KW. 1973. Routine haematological methods for use with fish blood. *J Fish Biol* 5 (4): 771-781. DOI: 10.1111/j.1095-8649.1973.tb04510.x.
- Borlongan LG. 1990. Studies on the digestive lipases of milkfish, *Chanos chanos*. *Aquaculture* 89 (1-3): 315-325. DOI: 10.1016/0044-8486(90)90135-A.
- Camevali O, Maradonna F, Gioacchini G. 2017. Integrated control of fish metabolism, wellbeing and reproduction: The role of probiotic. *Aquac* 472: 155-155. DOI: 10.1016/j.aquaculture.2016.03.037.
- Dai B, Hou Y, Hou Y, Qian L. 2018. Effects of multienzyme complex and probiotic supplementation on the growth performance, digestive enzyme activity and gut microorganisms composition of snakehead (*Channa argus*). *Aquac Nutr* 25 (1): 15-25. DOI: 10.1111/anu.12825.
- Darafsh F, Soltani M, Abdolhay HA, Mehrejan MS. 2019. Improvement of growth performance, digestive enzymes and body composition of Persian sturgeon (*Acipenser persicus*) following feeding on probiotic: *Bacillus licheniformis*, *Bacillus subtilis* and *Saccharomyces cerevisiae*. *Aquac Res* 51 (3): 957-964. DOI: 10.1111/are.14440.
- Dawood MAO, Koshio S, Ishikawa M, El-Sabagh M, Esteban AA, Zaineldin AI. 2016. Probiotics as an environment-friendly approach to enhance red sea bream, *Pagrus major* growth, immune response and oxidative status. *Fish Shellfish Immunol* 57: 170-178. DOI: 10.1016/j.fsi.2016.08.038.
- Dawood MAO, Koshio S. 2016. Recent advances in the role probiotics and prebiotics in carp aquaculture: A review. *Aquac* 454: 243-251. DOI: 10.1016/j.aquaculture.2015.12.03.
- Djauhari R, Widanarni, Sukenda, Suprayudi MA, Zairin Jr. M. 2016. Characterization of *Bacillus* sp. NP and its application as probiotic for common carp (*Cyprinus carpio*). *Res J Microbiol* 11 (4-5): 101-111. DOI: 10.3923/jm.2016.101.111.
- Doan HV, Doolgindachbaporn S, Suksri A. 2015. Effect of *LactoBacillus plantarum* and Jerusalem artichoke (*Helianthus tuberosus*) on growth performance, immunity and disease resistance of pangasius catfish (*Pangasius bocourti*, Sauvage 1880). *Aquac Nutr* 22 (2): 444-456. DOI: 10.1111/anu.12263.
- Elsabagh M, Mohamed R, Moustafa EM, Hamza A, Farrag F, Decamp O, Dawood MAO, Eltholth, M. 2018. Assessing the impact *Bacillus* strains mixture probiotic on water quality, growth performance, blood profile and intestinal morphology of Nile-Tilapia, *Oreochromis niloticus*. *Aquac Nutr* 24 (6): 1613-1622. DOI: 10.1111/anu.12797.
- Enferadi MHN, Mohammadizadeh F, Soltani M, Bahri AH, Sheikhzadeh N. 2018. Effects of *LactoBacillus plantarum* on growth performance, proteolytic enzymes activity and intestine morphology in rainbow trout (*Oncorhynchus mykiss*). *Turkish J Fisheries Aquat Sci* 18: 351-356. DOI: 10.4194/1303-2712-v18_2_14.
- Febrianti D, Yuhana M, Widanarni. 2016. Dietary symbiotic microcapsule influences the immune responses, growth performance and microbial population to white spot syndrome virus in pacific white shrimp (*Litopenaeus vannamei*). *J Fisheries Aquat Sci* 11: 28-42. DOI: 10.3923/jfas.2016.28.42.
- Food and Agriculture Organization of the United Nations (FAO). 2020. *The state of world fisheries and aquaculture 2020*. Sustainability in Action. Roma, Italia. DOI: 10.4060/ca9229en.
- Giri SS, Sen SS, Sukumaran V. 2012. Effects of dietary supplementation of potential probiotic *Pseudomonas aeruginosa* VSG-2 on the innate immunity and disease resistance of tropical freshwater fish, *Labeo rohita*. *Fish Shellfish Immunol* 32 (6): 1135-1140. DOI: 10.1016/j.fsi.2012.03.019.
- Hassani MHS, Jourdehi AY, Zelti AH, Masouleh AS, Lakani FB. 2019. Effects of commercial superzist probiotic on growth performance and hematological and immune indices in fingerlings *Acipenser baerii*. *Aquac Intl* 28: 377-387. DOI: 10.1007/s10499-019-00468-1.
- He R-P, Feng J, Tian X-L, Dong S-L, Wen B. 2017. Effects of dietary supplementation of probiotics on the growth, activities of digestive and non-specific immune enzymes in hybrid grouper (*Epinephelus lanceolatus* ♂ x *Epinephelus fuscoguttatus* ♀). *Aquac Res* 48 (12): 5782-5790. DOI: 10.1111/are.13401.
- Hoseiniyar SH, Sun Y-Z, Wang A, Zhou Z. 2018. Probiotic as means of diseases control in aquaculture, a review of current knowledge and future perspectives. *Front Microbiol* 9: 2429. DOI: 10.3389/fmicb.2018.02429.
- Huisman EA. 1987. *Principles of Fish Production*. Department of Fish Culture and Fisheries, Wageningen Agriculture University, Wageningen, Netherland.
- Ispir U, Dorucu M. 2005. A study on the effects of levamisole on immune system of Rainbow trout (*Oncorhynchus mykiss*, Walbaum). *Turkish J Vet Anim Sci* 29: 1169-1176.
- Jang WJ, Lee JM, Hasan TM, Lee B-J, Lim SG, Kong I-S. 2019. Effects of probiotic supplementation of a plant-based protein diet on intestinal microbial diversity, digestive enzyme activity, intestinal structure, and immunity in olive flounder (*Paralichthys olivaceus*). *Fish Shellfish Immunol* 92: 719-727. DOI: 10.1016/j.fsi.2019.06.056.

- Jung-Schroers V, Adamek M, Harris S, Syakuri H, Jung A, Irnazarow I, Steinhagen D. 2018. Response of the intestinal mucosal barrier of carp (*Cyprinus carpio*) to a bacterial challenge by *Aeromonas hydrophila* intubation after feeding with β -1,3/1,6-glucan. *J Fish Dis* 41 (7): 1077-1092. DOI: 10.1111/jfd.12799.
- Kewcharoen W, Srisapoom P. 2019. Probiotic effects of *Bacillus* spp. from Pacific white shrimp (*Litopenaeus vannamei*) on water quality and shrimp growth, immune responses and resistance to *Vibrio parahaemolyticus* (AHPND strains). *Fish Shellfish Immunol* 94: 175-189. DOI: 10.1016/j.fsi.2019.09.013.
- Komi FMM, Khalil F. 2017. Effect of ginger and ist nanoparticles on growth performance, cognition capability, immunity and prevention of motile *Aeromonas septicemia* in *Cyprinus carpio* fingerlings. *Aquac Nutr* 23 (6): 1492-1499. DOI: 10.1111/anu.12526.
- Kuebutornye FKA, Abarike ED, Lu Y. 2019. A review on the application of *Bacillus* as probiotic in aquaculture. *Fish Shellfish Immunol* 87: 820-828. DOI: 10.1016/j.fsi.2019.02.010.
- Kumar NR, Raman RP, Jadhao SJ, Brahmchari RK, Kumar K, Dash G. 2013. Effect of dietary supplementation of *Bacillus licheniformis* on gut microbiota, growth and immune response in giant freshwater prawn, *Macrobrachium rosenbergii* (de Man, 1879). *Aquac Intl* 21: 387-403. DOI: 10.1007/s10499-012-9567-8.
- Kumar R, Mukherjee SC, Ranjan R, Vani T, Brahmachari RK, Nayak SK. 2015. Effect of dietary supplementation of *Bacillus subtilis* on haematological and immunological parameters of *Catla catla* (Hamilton). *Aquac Intl* 23: 1275-1292. DOI: 10.1007/s10499-015-9883-x.
- Lee C, Cha J-H, Kim M-G, Shin J, Woo SH, Kim SH, Kim JW, Ji S-C, Lee K-J. 2019. The effects of dietary *Bacillus subtilis* on immune response, hematological parameters, growth performance and resistance of juvenile olive flounder (*Paralichthys olivaceus*) against *Streptococcus iniae*. *J World Aquac Soc* 51 (2): 551-562. DOI: 10.1111/jwas.12680.
- Lee W, Oh JY, Kim E-A, Kang N, Kim K-N, Ahn G, Jeon Y-J. 2016. A prebiotic role of *Ecklonia cava* improves the mortality of *Edwardsiella tarda*-infected zebrafish via regulating the growth of lactic acid bacteria and pathogen bacteria. *Fish Shellfish Immunol* 54: 620-628. DOI: 10.1016/j.fsi.2016.05.018.
- Li F, Wu D, Gu H-R, Yin M, Ge H-L, Liu X-H, Huang J, Zhang Y-G, Wang Z-J. 2019a. *Aeromonas hydrophila* and *Aeromonas veronii* cause motile *Aeromonas septicemia* in the cultured Chinese sucker, *Myxocyprinus asaticus*. *Aquac Res* 50 (5): 1515-1526. DOI: 10.1111/are.14028.
- Li J, Wu Z-B, Zhang Z, Zha J-W, Qu S-Y, Qi Q-Z, Wang G-X, Ling F. 2019b. Effects of potential probiotic *Bacillus velezensis* K2 on growth, immunity and resistance to *Vibrio harveyi* infection of hybrid grouper (*Epinephelus lanceolatus* ♂ x *Epinephelus fuscoguttatus* ♀). *Fish Shellfish Immunol* 93: 1047-1055. DOI: 10.1016/j.fsi.2019.08.047.
- Lin HZ, Guo Z, Yang Y, Zheng W, Li ZJ. 2004. Effect of dietary probiotics on apparent digestibility coefficients of nutrients of white shrimp *Litopenaeus vannamei* Boone. *Aquac Res* 35 (15): 1441-1447. DOI: 10.1111/j.1365-2109.2004.01169.x.
- Liu C-H, Wu K, Chu T-W, Wu T-M. 2017. Dietary supplementation of probiotic, *Bacillus subtilis* E20, enhances the growth performance and disease resistance against *Vibrio alginolyticus* in parrot fish (*Oplegnathus fasciatus*). *Aquac Intl* 26 (1): 63-74. DOI: 10.1007/s10499-017-0189-z.
- Madani NSH, Adorian TJ, Farsani HG, Hoseinifar SH. 2018. The effects of dietary probiotic Bacilli (*Bacillus subtilis* and *Bacillus licheniformis*) on growth performance, feed efficiency, body composition and immune parameters of white leg shrimp (*Litopenaeus vannamei*) postlarvae. *Aquac Res* 49 (5): 1926-1933. DOI: 10.1111/are.13648.
- Maji UJ, Mohanty S, Pradhan A, Maiti NK. 2017. Immune modulation, disease resistance and growth performance of Indian farmed carp, *Labeo rohita* (Hamilton), in response to dietary consortium of putative lactic acid bacteria. *Aquac Intl* 25: 1391-1407. DOI: 10.1007/s10499-017-0122-5.
- Ministry of Marine Affairs and Fisheries of the Republic Indonesia. 2019. Kelautan dan Perikanan dalam Angka. <https://knp.go.id/setjen/satudata/page/1453-kelautan-dan-perikanan-dalam-angka>. [Indonesian]
- Midhun SJ, Arun D, Sahadevan N, Vysakh A, Radhakrishnan EK, Jyothis M. 2019. Administration of probiotic *PaeniBacillus polymyxa* HGA4C induces morphometric, enzymatic and gene expression changes in *Oreochromis niloticus*. *Aquac* 508: 52-59. DOI: 10.1016/j.aquaculture.2019.04.061.
- Mohapatra S, Chakraborty T, Prusty AK, Das P, Paniprasad K, Mohanta KN. 2012. Use of different microbial probiotics in the diet of rohu, *Labeo rohita* fingerling: Effect on growth, nutrient digestibility and retention, digestive enzyme activities and intestinal microflora. *Aquac Nutr* 18 (1): 1-11. DOI: 10.1111/j.1365-2095.2011.00866.x.
- Nandi A, Banerjee G, Dan SK, Ghosh K, Ray AK. 2017. Probiotic efficiency of *Bacillus* sp. in *Labeo rohita* challenged by *Aeromonas hydrophila*: assessment of stress profile, haemato-biochemical parameters and immune responses. *Aquac Res* 48 (8): 4334-4345. DOI: 10.1111/are.13255.
- National Research Council (NRC). 1993. Nutrient Requirement of Fish. National Academy Press, Washington DC, USA.
- Nargesi EA, Falahatkar B, Sajjadi MM. 2019. Dietary supplementation of probiotic and influence on feed efficiency, growth parameters and reproductive performance in female rainbow trout (*Oncorhynchus mykiss*) broodstock. *Aquac Nutr* 26 (1): 98-108. DOI: 10.1111/anu.12970.
- Olivier G, Lallier R, Lariviere S. 1981. A toxigenic profile of *Aeromonas hydrophila* and *Aeromonas sobria* isolated from fish. *Canadian J Microbiol* 27 (3): 330-333. DOI: 10.1139/m81-050.
- Poolsawat L, Li X, He M, Ji D, Leng X. 2019. *Clostridium butyricum* as probiotic for promoting growth performance, feed utilization gut health and microbiota community of tilapia (*Oreochromis niloticus* x *O. aureus*). *Aquac Nutr* 26 (3): 657-670. DOI: 10.1111/anu.13025.
- Pourgholam MA, Khara H, Safari R, Sadati MAY, Aramli MS. 2017. Influence of *Lactobacillus plantarum* inclusion in the diet of Siberian sturgeon (*Acipenser baeri*) on performance and hematological parameters. *Turkish J Fisheries Aquat Sci* 17: 1-5. DOI: 10.4194/1303-2712-v17_1_01.
- Putra AN, Widanarni. 2015. Screening of amylolytic bacteria as candidates of probiotic in tilapia *Oreochromis* sp. *Res J Microbiol* 10: 1-13. DOI: 10.3923/jm.2015.1.13.
- Putra AN, Widanarni, Utomo NBP. 2015. Growth performance of tilapia (*Oreochromis niloticus*) fed with probiotic, prebiotic and symbiotic in diet. *Pak J Nutr* 14: 263-268. DOI: 10.3923/pjn.2015.263.268.
- Putra AN, Romdhonah Y. 2019. Effects of dietary *Bacillus* NP5 and sweet potato extract on growth and digestive enzyme activity of dumbo catfish, *Clarias* sp.. *Jurnal Akuakultur Indonesia* 18 (1): 80-88. DOI: 10.19027/jai.18.1.80-88.
- Putra AN, Syamsunamo MB, Ningrum W, Jumyanah, Mustahal. 2020a. Effect of the administration of probiotic *Bacillus* NP5 in the rearing media on water quality, growth, and disease resistance of African catfish (*Clarias gariepinus*). *Biodiversitas* 21 (6): 2566-2575. DOI: 10.13057/biodiv/d210629.
- Putra AN, Mustahal, Syamsunamo MB. 2020b. Effects of dietary probiotic *Bacillus* NP5 on the growth performances of catfish (*Clarias* sp.). *Biotropia* 27 (1): 51-59. DOI: 10.11598/btb.2020.27.1.1102.
- Ramesh D, Vinothkanna A, Rai AK, Vignesh VS. 2015. Isolation of potential probiotic *Bacillus* spp. and assessment of their subcellular components to induce immune responses in *Labeo rohita* against *Aeromonas hydrophila*. *Fish Shellfish Immunol* 45: 268-276. DOI: 10.1016/j.fsi.2015.04.018.
- Reda RM, Selim KM. 2015. Evaluation of *Bacillus amyloliquefaciens* on the growth performance, intestinal morphology, hematology and body composition of Nile tilapia, *Oreochromis niloticus*. *Aquac Intl* 23: 203-217. DOI: 10.1007/s10499-014-9809-z.
- RingØ E. 2020. Probiotics in shellfish aquaculture. *Aquac Fisheries* 5: 1-27. DOI: 10.1016/j.aaf.2019.12.001.
- Sankar H, Philip B, Philip R, Singh ISB. 2016. Effect of probiotic on digestive enzyme activities and growth of cichlids, *Etroplus suratensis* (Pearl spot) and *Oreochromis mossambicus* (Tilapia). *Aquac Nutr* 23 (4): 852-864. DOI: 10.1111/anu.12452.
- Shoemaker CA, Mohammed HH, Bader TJ, Peatman E, Beck BH. 2018. Immersion vaccination with an inactivated virulent *Aeromonas hydrophila* bacterin protects hybrid catfish (*Ictalurus punctatus* X *Ictalurus furcatus*) from motile *Aeromonas septicemia*. *Fish Shellfish Immunol* 82: 239-242. DOI: 10.1016/j.fsi.2018.08.040.
- Soltani M, Abdy E, Alishahi M, Mirghaed AT, Hosseini-Shekarabi P. 2017. Growth performance, immune-physiological variables and disease resistance of common carp (*Cyprinus carpio*) orally subjected to different concentrations of *LactoBacillus plantarum*. *Aquac Intl* 25: 1913-1933. DOI: 10.1007/s10499-017-0164-8.

- Takeuchi T. 1988. Laboratory work-chemical evaluation of dietary nutrients. In: Watanabe T (eds). Fish Nutrition and Mariculture. Kanagawa International Fisheries Training Centre, Japan International Cooperation Agency (JICA), Japan.
- Tamamduhuri R, Widanarni, Yuhana M. 2016. Administration of microencapsulated probiotic *Bacillus* sp. NP5 and prebiotic mannan oligosaccharide for prevention of *Aeromonas hydrophila* infection on *Pangasianodon hypophthalmus*. J Fisheries Aquat Sci 11: 67-76. DOI: 10.3923/jfas.2016.67.76.
- Tanbiyaskur, Widanarni, Lusiatuti AM. 2015. Administration of *Bacillus* NP5 and oligosaccharide to enhance the immune response in tilapia *Oreochromis niloticus* towards streptococcosis. Intl J Sci Basic Appl Res 20 (2): 301-315.
- Tantikitti C. 2014. Review article: Feed palatability and the alternative protein sources in shrimp feed. Songklanakarini J Sci Technol 36 (1): 51-55.
- Tsai C-Y, Chi C-C, Liu C-H. 2019. The growth and apparent digestibility of white shrimp, *Litopenaeus vannamei*, are increased with the probiotic, *Bacillus subtilis*. Aquac Res 50 (5): 1475-1481. DOI: 10.1111/are.14022.
- Uribe C, Folch H, Enriquez R, Moran. 2011. Innate and adaptive immunity in teleost fish: A review. Veterinarni Medicina 56 (10): 486-503. DOI: 10.17221/3294-VETMED.
- Utami DAS, Widanarni, Suprayudi MA. 2015. Quality of dried *Bacillus* NP5 and its effect on growth performance of tilapia *Oreochromis niloticus*. Pak J Biol Sci 18: 88-93. DOI: 10.3923/pjbs.2015.88.93.
- Valiallahi J, Pouranasali M, Janalizadeh E, Bucio A. 2018. Use of *LactoBacillus* for improves growth and enhanced biochemical, hematological, and digestive enzyme activity in common carp at Mazandaran, Iran. North Am J Aquac 80 (2): 206-215. DOI: 10.1002/naaq.10027.
- Wedemeyer GA, Yasutake. 1977. Clinical Methods for the Assessment on the Effect of Environmental Stress on Fish Health. US Fish and Wildlife Service, USA.
- Widanarni, Yuhana M, Muhammad A. 2014. *Bacillus* NP5 improves growth performance and resistance against infectious myonecrosis virus in white shrimp *Litopenaeus vannamei*. Ilmu Kelautan 19: 211-218. DOI: 10.14710/ik.ijms.19.4.211-218.
- Worthington V. 1993. Worthington Enzyme Manual. Enzymes and Related Biochemicals. Worthington Chemical Corp, USA.
- Wu ZX, Feng X, Xie LL, Peng XY, Chen XX. 2012. Effect of probiotic *Bacillus subtilis* Ch9 for grass carp, *Ctenopharyngodon idella* (Valenciennes, 1844), on growth performance, digestive enzyme activities and intestinal microflora. J Appl Ichthyol 28 (5): 721-727. DOI: 10.1111/j.1439-0426.2012.01968.x.
- Xie J-J, Liu Q-Q, Liao S, Fang H-H, Yin P, Xie S-W, Tian X, Liu Y-J, Niu J. 2019. Effects of dietary mixed probiotics on growth, non-specific immunity, intestinal morphology and microbiota of juvenile pacific white shrimp, *Litopenaeus vannamei*. Fish Shellfish Immunol 90: 456-465. DOI: 10.1016/j.fsi.2019.04.301.
- Zhang C-N, Zhang J-L, Guan W-C, Zhang X-F, Guan S-H, Zeng Q-H, Cheng G-F, Cui W. 2017. Effects of *Lactobacillus delbrueckii* on immune response, disease resistance against *Aeromonas hydrophila*, antioxidant capability and growth performance of *Cyprinus carpio* Huanghe var. Fish Shellfish Immunol 68: 84-91. DOI: 10.1016/j.fsi.2017.07.012.
- Zhang C, Zhang J, Fan W, Huang M, Liu M. 2018. Effects of dietary *Lactobacillus delbrueckii* on growth performance, body composition, digestive and absorptive capacity, and gene expression of common carp (*Cyprinus carpio* Huanghe var). Aquac Nutr 25 (1): 166-175. DOI: 10.1111/anu.12840.
- Zhao C, Zhu J, Hu J, Dong X, Sun L, Zhang X, Miao S. 2019. Effects of dietary *Bacillus pumilus* on growth performance, innate immunity and digestive enzymes of giant freshwater prawns (*Macrobrachium rosenbergii*). Aquac Nutr 25 (3): 712-720. DOI: 10.1111/anu.12894.
- Zheng CN, Wang W. 2016. Effects of *Lactobacillus pentosus* on the growth performance, digestive enzyme and disease resistance of white shrimp, *Litopenaeus vannamei* (Boone, 1931). Aquac Res 48 (6): 2767-2777. DOI: 10.1111/are.13110.
- Zhu C, Yu L, Liu W, Jiang M, He S, Yi G, Wen H, Liang X. 2019. Dietary supplementation with *Bacillus subtilis* LT3-3 enhances the growth, immunity and disease resistance against *Streptococcus agalactiae* infection in genetically improved farmed tilapia, *Oreochromis niloticus*. Aquac Nutr 25 (6): 1241-1249. DOI: 10.1111/anu.12938.
- Zokaeifar H, Balcazar JL, Saad CR, Kamarudin MS, Sijam K, Arshad A, Nejat N. 2012. Effect of *Bacillus subtilis* on the growth performance, digestive enzymes, immune gene expression and disease resistance of white shrimp, *Litopenaeus vannamei*. Fish Shellfish Immunol 33: 683-689. DOI: 10.1016/j.fsi.2012.05.027.
- Zokaeifar H, Babaei N, Saad CR, Kamarudin MS, Sijam K, Balcazar JL. 2014. Administration of *Bacillus subtilis* strains in the rearing water enhances the water quality, growth performance, immune response, and resistance against *Vibrio harveyi* infection in juvenile white shrimp, *Litopenaeus vannamei*. Fish Shellfish Immunol 36 (1): 68-74. DOI: 10.1016/j.fsi.2013.10.007.

biodiversitas 2021

ORIGINALITY REPORT

19%

SIMILARITY INDEX

14%

INTERNET SOURCES

14%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1	journal.ipb.ac.id Internet Source	1%
2	www.bioflux.com.ro Internet Source	1%
3	www.tandfonline.com Internet Source	1%
4	tailieu.vn Internet Source	1%
5	etda.libraries.psu.edu Internet Source	1%
6	www.omicsonline.org Internet Source	1%
7	"Probiotic Bacteria and Postbiotic Metabolites: Role in Animal and Human Health", Springer Science and Business Media LLC, 2021 Publication	<1%
8	repository.untad.ac.id Internet Source	<1%

9	www.mdpi.com Internet Source	<1 %
10	ejfa.me Internet Source	<1 %
11	www.fisheriesjournal.com Internet Source	<1 %
12	Brian Austin, Dawn A. Austin. "Bacterial Fish Pathogens", Springer Science and Business Media LLC, 2016 Publication	<1 %
13	Reza Tarkhani, Ahmad Imani, Seyed Hossein Hoseinifar, Kourosch Sarvi Moghanlou, Ramin Manaffar. "The effects of host-associated <i>Enterococcus faecium</i> CGMCC1.2136 on serum immune parameters, digestive enzymes activity and growth performance of the Caspian roach (<i>Rutilus rutilus caspicus</i>) fingerlings", <i>Aquaculture</i> , 2020 Publication	<1 %
14	archimer.ifremer.fr Internet Source	<1 %
15	www.ncbi.nlm.nih.gov Internet Source	<1 %
16	Lukman Anugrah Agung, Widanarni ., Munti Yuhana. "Application of Micro-Encapsulated Probiotic <i>Bacillus</i> NP5 and Prebiotic Mannan	<1 %

Oligosaccharide (MOS) to Prevent Streptococcosis on Tilapia Oreochromis niloticus", Research Journal of Microbiology, 2015

Publication

17

mafiadoc.com

Internet Source

<1 %

18

Daniel Assan, Felix K.A. Kuebutornye, Vivian Hlordzi, Huapu Chen, Jan Mraz, Umar Farouk Mustapha, Emmanuel D. Abarike. "Effects of probiotics on digestive enzymes of fish (finfish and shellfish); status and prospects: A mini review", Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology, 2021

Publication

<1 %

19

opus.uni-hohenheim.de

Internet Source

<1 %

20

Greeshma James, Bini C. Das, Seena Jose, Rejish Kumar V.J. "Bacillus as an aquaculture friendly microbe", Aquaculture International, 2021

Publication

<1 %

21

Hien Van Doan, Seyed Hossein Hoseinifar, Wanaporn Tapingkae, Mintra Seel-audom et al. "Boosted Growth Performance, Mucosal and Serum Immunity, and Disease Resistance Nile Tilapia (Oreochromis niloticus) Fingerlings

<1 %

Using Corncob-Derived Xylooligosaccharide and Lactobacillus plantarum CR1T5",
Probiotics and Antimicrobial Proteins, 2019

Publication

22

www.science.gov

Internet Source

<1 %

23

Govindaraj Krishnaveni, Samayanpaulraj Vignesh, Narayanadoss Vidhyalakshmi, Velu Vijay, Uthandakalaipandian Ramesh. " Effects of dietary supplementation of URLP18 on growth, innate immunity and survival against ATCC 7966 challenge in freshwater fish (common carp) ", Aquaculture Research, 2020

Publication

<1 %

24

Darío Israel García-Medel, Carlos Angulo, Ruth Escamilla-Montes, Jesús Arturo Fierro-Coronado et al. "Bacillus licheniformis BCR 4-3 increases immune response and survival of Litopenaeus vannamei challenged with Vibrio parahaemolyticus IPNGS16", Aquaculture International, 2020

Publication

<1 %

25

Wenxiang Yao, Xiaoqin Li, Chunyan Zhang, Jing Wang, Youwang Cai, Xiangjun Leng. "Effects of dietary synbiotics supplementation methods on growth, intestinal health, non-specific immunity and disease resistance of

<1 %

Pacific white shrimp, *Litopenaeus vannamei*",
Fish & Shellfish Immunology, 2021

Publication

26

eprints.cmfri.org.in

Internet Source

<1 %

27

journals-crea.4science.it

Internet Source

<1 %

28

journals.tubitak.gov.tr

Internet Source

<1 %

29

Gang Yang, Kaikai Shen, Ruihe Yu, Qiong Wu, Quangen Yan, Wenjing Chen, Liyun Ding, Vikas Kumar, Chungen Wen, Mo Peng. "Probiotic (*Bacillus cereus*) enhanced growth of Pengze crucian carp concurrent with modulating the antioxidant defense response and exerting beneficial impacts on inflammatory response via Nrf2 activation", *Aquaculture*, 2020

Publication

<1 %

30

Jalal Valiollahi, Mohsen Pourabasali, Elham Janalizadeh, Adolfo Bucio. " Use of for Improved Growth and Enhanced Biochemical, Hematological, and Digestive Enzyme Activity in Common Carp at Mazandaran, Iran ", *North American Journal of Aquaculture*, 2018

Publication

<1 %

31

Rajabdeen Jannathulla, Oduru Sravanthi, Syed Moomeen, Gopalapillai Gopikrishna,

<1 %

Jagabattula Syama Dayal. "Microbial products in terms of isolates, whole-cell biomass, and live organisms as aquafeed ingredients: production, nutritional values, and market potential—a review", *Aquaculture International*, 2021

Publication

32

Yalda Hooshyar, Abdolmohammad Abedian Kenari, Hamed Paknejad, Hassan Gandomi. "Effects of *Lactobacillus rhamnosus* ATCC 7469 on Different Parameters Related to Health Status of Rainbow Trout (*Oncorhynchus mykiss*) and the Protection Against *Yersinia ruckeri*", *Probiotics and Antimicrobial Proteins*, 2020

Publication

<1 %

33

cwww.intechopen.com

Internet Source

<1 %

34

eprints.keele.ac.uk

Internet Source

<1 %

35

fendergisi.selcuk.edu.tr

Internet Source

<1 %

36

munin.uit.no

Internet Source

<1 %

37

onlinelibrary.wiley.com

Internet Source

<1 %

38

www.icddrb.org

Internet Source

<1 %

39

www.pjbs.org

Internet Source

<1 %

40

www.yumpu.com

Internet Source

<1 %

41

Gang Yang, Hongzhong Cao, Wenhao Jiang, Baoqing Hu, Shaoqing Jian, Chungen Wen, Kimia Kajbaf, Vikas Kumar, Zhiying Tao, Mo Peng. " Dietary supplementation of as probiotics in Pengze crucian carp (var. Pengze): Effects on growth performance, fillet quality, serum biochemical parameters and intestinal histology ", Aquaculture Research, 2019

Publication

<1 %

42

Hooshang Yeganeh Rastekenari, Rezvanollah Kazami, Alireza Shenavar Masouleh, Akbar Banavreh et al. " Autochthonous probiotics and in the diet of fingerlings great sturgeon, : effects on growth performance, feed efficiency, haematological parameters, immune status and intestinal morphology ", Aquaculture Research, 2021

Publication

<1 %

43

Md. Mahbubur Rahman, A. Q. M. Robiul Kawser, Md Tofazzal Islam. "Chapter 16

<1 %

Probiotic Bacilli in Sustainable Aquaculture",
Springer Science and Business Media LLC,
2019

Publication

44

Sakineh Yeganeh, Milad Adel, Ahmad Nosratimovafagh, Mahmoud A. O. Dawood. "The Effect of Lactococcus lactis subsp. lactis PTCC 1403 on the Growth Performance, Digestive Enzymes Activity, Antioxidative Status, Immune Response, and Disease Resistance of Rainbow Trout (*Oncorhynchus mykiss*)", Probiotics and Antimicrobial Proteins, 2021

Publication

<1 %

45

Sib Sankar Giri, Jin Woo Jun, Saekil Yun, Hyoun Joong Kim et al. "Effects of dietary heat-killed *Pseudomonas aeruginosa* strain VSG2 on immune functions, antioxidant efficacy, and disease resistance in *Cyprinus carpio*", Aquaculture, 2020

Publication

<1 %

46

ccrjournal.com
Internet Source

<1 %

47

garuda.ristekbrin.go.id
Internet Source

<1 %

48

journal.ke.hu
Internet Source

<1 %

49

ntnuopen.ntnu.no

Internet Source

<1 %

50

www.frontiersin.org

Internet Source

<1 %

51

Dawood, Mahmoud A.O., Shunsuke Koshio, Mabrouk El-Sabagh, Md Masum Billah, Amr I. Zaineldin, Mohamed Mamdouh Zayed, and Amira Alaa El-Dein Omar. "Changes in the growth, humoral and mucosal immune responses following β -glucan and vitamin C administration in red sea bream, *Pagrus major*", *Aquaculture*, 2016.

Publication

<1 %

52

Defeng Zhang, Yanxia Gao, Xiaoli Ke, Mengmeng Yi, Zhigang Liu, Xueqing Han, Cunbin Shi, Maixin Lu. "Bacillus velezensis LF01: in vitro antimicrobial activity against fish pathogens, growth performance enhancement, and disease resistance against streptococcosis in Nile tilapia (*Oreochromis niloticus*)", *Applied Microbiology and Biotechnology*, 2019

Publication

<1 %

53

Dharmaraj Ramesh, Sami Souissi. " Effects of potential probiotic KADR1 and its subcellular components on immune responses and

<1 %

disease resistance in ", Aquaculture Research, 2018

Publication

54

Md. Idrish Raja Khan, Dibyendu Kamilya, Tanmoy Gon Choudhury, Partha Sarathi Tripathy, Gaurav Rathore. "Deciphering the Probiotic Potential of *Bacillus amyloliquefaciens* COFCAU_P1 Isolated from the Intestine of *Labeo rohita* Through In Vitro and Genetic Assessment", *Probiotics and Antimicrobial Proteins*, 2021

Publication

<1 %

55

Mehdi Soltani, Esmail Abdy, Mojtaba Alishahi, Ali Taheri Mirghaed, Pezhman Hosseini-Shekarabi. "Growth performance, immune-physiological variables and disease resistance of common carp (*Cyprinus carpio*) orally subjected to different concentrations of *Lactobacillus plantarum*", *Aquaculture International*, 2017

Publication

<1 %

56

Ngoc Tuan Tran, Zhongzhen Li, Hongyu Ma, Yueling Zhang, Huaiping Zheng, Yi Gong, Shengkang Li. " : a promising probiotic confers positive health benefits in aquatic animals ", *Reviews in Aquaculture*, 2020

Publication

<1 %

57 Nuno Borges, Tina Keller-Costa, Gracinda M.M. Sanches-Fernandes, António Louvado, Newton C.M. Gomes, Rodrigo Costa. "Bacteriome Structure, Function, and Probiotics in Fish Larviculture: The Good, the Bad, and the Gaps", Annual Review of Animal Biosciences, 2021
Publication

58 boku.ac.at
Internet Source

59 doaj.org
Internet Source

60 ejournal-balitbang.kkp.go.id
Internet Source

61 ir.kagoshima-u.ac.jp
Internet Source

62 lppm.ub.ac.id
Internet Source

63 ojs.omniakuatika.net
Internet Source

64 repositorio-aberto.up.pt
Internet Source

65 www.ijsrp.org
Internet Source

66 www.scribd.com

<1 %

67

www.trnres.com

Internet Source

<1 %

68

Chorong Lee, Ji - Hoon Cha, Min - Gi Kim, Jaehyeong Shin et al. " The effects of dietary on immune response, hematological parameters, growth performance, and resistance of juvenile olive flounder () against ", Journal of the World Aquaculture Society, 2019

Publication

<1 %

69

A N Putra, Jaenudin, F Sofia, Mustahal, M B Syamsunarno, D Hermawan, M Herjayanto. " The Utilization of Vegetable Waste Silage as Feed Ingredient in Diets for Tilapia ", IOP Conference Series: Earth and Environmental Science, 2019

Publication

<1 %

70

Bidhan C. De, D. K. Meena, B. K. Behera, Pronob Das, P. K. Das Mohapatra, A. P. Sharma. "Probiotics in fish and shellfish culture: immunomodulatory and ecophysiological responses", Fish Physiology and Biochemistry, 2014

Publication

<1 %

71

Hien Van Doan, Mehdi Soltani, Einar Ringø. "In vitro antagonistic effect and in vivo protective efficacy of Gram-positive probiotics versus Gram-negative bacterial pathogens in finfish and shellfish", Aquaculture, 2021

Publication

<1 %

72

Nisha Chuphal, Krishna Pada Singha, Parimal Sardar, Narottam Prasad Sahu, Naseemashahul Shamna, Vikas Kumar. "Scope of Archaea in Fish Feed: a New Chapter in Aquafeed Probiotics?", Probiotics and Antimicrobial Proteins, 2021

Publication

<1 %

73

Ricky Djauhari, Widanarni ., Sukenda ., Muhammad Agus Suprayudi, Muhammad Zairin Jr.. "Characterization of Bacillus sp. NP5 and its Application as Probiotic for Common Carp (Cyprinus carpio)", Research Journal of Microbiology, 2016

Publication

<1 %

74

Rifqi Tamamdustu, Widanarni ., Munti Yuhana. "Administration of Microencapsulated Probiotic Bacillus sp. NP5 and Prebiotic Mannan Oligosaccharide for Prevention of Aeromonas hydrophila Infection on Pangasianodon hypophthalmus", Journal of Fisheries and Aquatic Science, 2016

Publication

<1 %

75

Seyed Hossein Hoseinifar, Maryam Dadar, Einar Ringø. "Modulation of nutrient digestibility and digestive enzyme activities in aquatic animals: The functional feed additives scenario", *Aquaculture Research*, 2017

Publication

<1 %

76

Vivian Hlordzi, Felix K.A. Kuebutornye, Gyamfua Afriyie, Emmanuel Delwin Abarike, Yishan Lu, Shuyan Chi, Melody A. Anokyewaa. "The use of *Bacillus* species in maintenance of water quality in aquaculture: A review", *Aquaculture Reports*, 2020

Publication

<1 %

77

Yan Cai, Wei Yuan, Shifeng Wang, Weiliang Guo, An Li, Yue Wu, Xin Chen, Zhuling Ren, Yongcan Zhou. "In vitro screening of putative probiotics and their dual beneficial effects: To white shrimp (*Litopenaeus vannamei*) postlarvae and to the rearing water", *Aquaculture*, 2018

Publication

<1 %

78

Zequn Li, Ning Bao, Tongjun Ren, Yuzhe Han, Zhiqiang Jiang, Zhuoan Bai, Yanan Hu, Jingyun Ding. "The effect of a multi-strain probiotic on growth performance, non-specific immune response, and intestinal health of juvenile turbot, *Scophthalmus maximus* L.", *Fish Physiology and Biochemistry*, 2019

<1 %

Publication

Exclude quotes On

Exclude bibliography On

Exclude matches Off

biodiversitas 2021

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9
